

## Referenties behorende bij de Multidisciplinaire Richtlijn Hartrevalidatie 2011

- Abbas AE et al. Frequency of returning to work one and six months following percutaneous coronary intervention for acute myocardial infarction. *Am J Cardiol* 2004;94(11):1403-5
- Adamopoulos S et al., Circadian pattern of heart rate variability in chronic heart failure patients. Effects of physical training. *Eur Heart J* 1995;16(10):1380-86
- Adamopoulos S, Coats AJ, Brunotte F, et al. Physical training improves skeletal muscle metabolism in patients with chronic heart failure. *J Am Coll Cardiol* 1993;21:1101-6
- Adamopoulos S, Parissis J, Karatzas D, et al. Physical training modulates proinflammatory cytokines and the soluble Fas/soluble Fas ligand system in patients with chronic heart failure. *J Am Coll Cardiol* 2002;39:653-63
- Adamopoulos S, Parissis J, Kroupis C, et al. Physical training reduces peripheral markers of inflammation in patients with chronic heart failure. *Eur Heart J* 2001;22:791-7
- Adamopoulos S, Ponikowski P, Cerquetani E, et al. Circadian pattern of heart rate variability in chronic heart failure patients. Effects of physical training. *Eur Heart J* 1995;16:1380-6
- Adams KJ, Barnard KL, Swank AM, et al. Combined high intensity strength and aerobic training in diverse phase II cardiac rehabilitation patients. *J Cardiopulm Rehabil* 1999;4:209-15
- Ades P, Pashkow F, Nestor J, et al. Cost-effectiveness of cardiac rehabilitation after myocardial infarction. *J Cardiopulm Rehabil* 1997;17:222-31
- Ades PA et al. Referral patterns and exercise response in the rehabilitation of female coronary patients aged greater than or equal to 62 years. *Am J Cardiol* 1992;69(17): 422-25.
- Ades PA, Malonney A, Savage P, et al. Determinants of physical functioning in coronary patients: response to cardiac rehabilitation. *Arch Intern Med* 1999;159:2357-60
- Ades PA. Cardiac rehabilitation and secondary prevention of coronary heart disease. *N Engl J Med* 2001;345(12):892-902
- Agren JJ, Vaisanen S, Hanninen O, et al. Hemostatic factors and platelet aggregation after a fish-enriched diet or fish oil or docosahexaenoic acid supplementation. *Prostaglandins Leukot Essent Fatty Acids* 1997;57:419-21
- Agren S et al. Addressing spouses' unique needs after cardiac surgery when recovery is complicated by heart failure. *Heart Lung* 2009;38(4):284-91
- Ala-Mursula L, Vahtera J, Linna A, et al. Employee worktime control moderates the effects of job strain and effort-reward imbalance on sickness absence: the 10-town study. *J Epidemiol Community Health* 2005;59(10):851-7
- Albert C, Mittleman M, Chae C. Triggering of sudden death by vigorous exertion. *N Engl J Med* 2000;343:1355-61
- Amarosa-Tuppler B, Tapp J, et al. Stress management through relaxation and imagery in the treatment of angina pectoris. *J Cardiac Rehabil* 1989;9:348-53

- Amin, A.A., et al., The prevalence of unrecognized depression in patients with acute coronary syndrome. *Am Heart J* 2006;152(5): 928-34
- Andre-Petersson L, Engstrom G, Hedblad B, et al. Social support at work and the risk of myocardial infarction and stroke in women and men. *Soc Sci Med* 2007;64(4):830-4.
- Anker SD et al., Hormonal changes and catabolic/anabolic imbalance in chronic heart failure and their importance for cardiac cachexia. *Circulation* 1997;96(2):526-34
- Antman EM, Anbe DT, Armstrong PW, et al. ACC/AHA guidelines for the management of patients with ST-elevation myocardial infarction: A report of the American College of Cardiology/American Heart Association Task Force on Practice Guidelines (committee to revise the 1999 guidelines for the management of patients with acute myocardial infarction). *J Am Coll Cardiol* 2004;44:E1-E211.
- Antonios TF, MacGregor GA. Salt-more adverse effects. *Lancet* 1996;348(9022): 250-51.
- Appel LJ, Champagne CM, Harsha DW, et al. Effects of comprehensive lifestyle modification on blood pressure control: main results of the PREMIER clinical trial. *JAMA* 2003;289:2083-93
- Appel LJ, Miller ER, Jee SH, et al. Effect of dietary patterns on serum homocysteine: results of a randomized, controlled feeding study. *Circulation* 2000;102:852-7
- Appel LJ, Moore TJ, Obarzanek E, et al. A clinical trial of the effects of dietary patterns on blood pressure. DASH Collaborative Research Group. *N Engl J Med* 1997;336:1117-24
- Appels A et al. Effects of a behavioural intervention on quality of life and related variables in angioplasty patients: results of the Exhaustion Intervention Trial. *J Psychosom Res* 2006;61(1):1-7
- Appels A et al., Effects of treating exhaustion in angioplasty patients on new coronary events: results of the randomized Exhaustion Intervention Trial (EXIT). *Psychosom Med*, 2005;67(2):217-23
- Appels A, Bär F, van der Pol G, et al. Effects of treating exhaustion in angioplasty patients on new coronary events: results of the randomized Exhaustion Intervention Trial (EXIT). *Psychosom Med* 2005;67(2):217-23
- Appels A, Bar FWH, Lasker J, et al. The effect of a psychological intervention program on the risk of a new coronary event after angioplasty: a feasibility study. *J Psychosom Res* 1997;17:209-17
- Appels A. Exhaustion and coronary heart disease: the history of a scientific quest. *Patient Educ Couns* 2004;55(2):223-9
- Arawwawala D, Brett SJ. Clinical review: beyond immediate survival from resuscitation-long-term outcome considerations after cardiac arrest. *Crit Care* 2007;11(6):235
- Arnold SV, Alexander KP, Masoudi FA, et al. The effect of age on functional and mortality outcomes after acute myocardial infarction. *J Am Geriatr Soc* 2009;57(2):209-17

- Arthur HM, Daniels C, McKelvie R, et al. Effect of a preoperative intervention on preoperative and postoperative outcomes in low-risk patients awaiting elective coronary artery bypass graft surgery. A randomized, controlled trial. *Ann Intern Med*. 2000;133(4):253-62
- Baars HMJ. Handleiding Maastrichtse Sociale Netwerk Analyse. In: Sociale netwerkstudies, cahier 16. Maastricht: Vakgroep Psychiatrie & Psychiatrische Epidemiologie, Universiteit Maastricht 1997,.
- Badenhop DT. The therapeutic role of exercise in patients with orthotopic heart transplant. *Med Sci Sports Exerc* 1995;27(7):975-85.
- Baer PE, Cleveland SE, Montero AC. Improving post-myocardial infarction recovery status by stress management training during hospitalization. *J Cardiopulm Rehabil* 1985; 5:191-6
- Baessler A, Hengstenberg C, Holmer S, et al. Long-term effects of in-hospital cardiac rehabilitation on the cardiac risk profile. *Eur Heart J* 2001;22(13):1111-8
- Bakas T et al. Family caregiving in heart failure. *Nurs Res*, 2006. 55(3): 180-8.
- Balestrieri M et al. Identification of depression by medical and surgical general hospital physicians. *Gen Hosp Psychiatry* 2002;24(1):4-11.
- Bankier B, Januzzi JL, Littman AB. The high prevalence of multiple psychiatric disorders in stable outpatients with coronary heart disease. *Psychosom Med* 2004;66(5):645-50
- Barnard ND, Akhtar A, Nicholson A. Factors that facilitate compliance to lower fat intake. *Arch Fam Med* 1995;4:153-8
- Bar-On D, Gilutz H, Maymon T, et al. Long-term prognosis of low-risk, post-MI patients: the importance of subjective perception of disease. *Eur Heart J*. 1994;15(12):1611-15
- Bar-On D. Causal attributions and the rehabilitation of myocardial infarction victims. *J Social Clin Psychol* 1987;5(1):114-22
- Barth J, Schumacher M, Herrmann-Lingen C. Depression as a risk factor for mortality in patients with coronary heart disease: a meta-analysis. *Psychosom Med* 2004;66(6):802-13
- Barzi F, Woodward M, Marfisi RM, et al. Mediterranean diet and all-causes mortality after myocardial infarction: results from the GISSI-Prevenzione trial. *Eur J Clin Nutr* 2003;57:604-11
- Bazzano LA, He J, Ogden LG, et al. Fruit and vegetable intake and risk of cardiovascular disease in US adults: the first National Health and Nutrition Examination Survey Epidemiologic Follow-up Study. *Am J Clin Nutr* 2002;76:93-9
- Belardinelli R, Georgiou D, Cianci G, et al. Randomized, controlled trial of long-term moderate exercise training in chronic heart failure: effects on functional capacity, quality of life, and clinical outcome. *Circulation* 1999;99:1173-82
- Belkic KL, Landsbergis PA, Schnall PL, Baker D. Is job strain a major source of cardiovascular disease risk? *Scand J Work Environ Health* 2004;30(2):85-128

- Bell JM. A comparison of a multidisciplinary home based cardiac rehabilitation programme with comprehensive conventional rehabilitation in post-myocardial infarction patients [PhD thesis]. London: University of London 1998
- Bemelmans WJ, Broer J, Feskens EJ, et al. Effect of an increased intake of alpha-linolenic acid and group nutritional education on cardiovascular risk factors: the Mediterranean Alpha-linolenic Enriched Groningen Dietary Intervention (MARGARIN) study. *Am J Clin Nutr* 2002;75:221-7
- Bengtsson K. Rehabilitation after myocardial infarction: a controlled study. *Scand J Rehabil Med* 1983;15(1):1-9
- Beniamini Y, Rubenstein J, Faigenbaum A, et al. High-intensity strength training of patients enrolled in an outpatient cardiac rehabilitation program. *J Cardiopulm Rehabil* 1999;19:8-17
- Beniaminovitz A, Lang CC, LaManca J, et al. Selective low-level leg muscle training alleviates dyspnea in patients with heart failure. *J Am Coll Cardiol* 2002;40:1602-8
- Berkman, L.F., et al., Effects of treating depression and low perceived social support on clinical events after myocardial infarction: the Enhancing Recovery in Coronary Heart Disease Patients (ENRICHED) Randomized Trial. *JAMA* 2003;289(23):3106-16
- Berlin JA, Colditz GA. A meta-analysis of physical activity in the prevention of coronary heart disease. *Am J Epidemiol* 1990;132:612-28
- Bethell HJ. Cardiac rehabilitation: from Hellerstein to the millennium. *Int J Clin Pract* 2000;54:92-7
- Bhattacharyya MR, Perkins-Porras L, Whitehead DL, et al. Psychological and clinical predictors of return to work after acute coronary syndrome. *Eur Heart J* 2007;28(2):160-65.
- Blumenthal JA et al. Exercise, depression, and mortality after myocardial infarction in the ENRICHED trial. *Med Sci Sports Exerc*, 2004. 36(5): 746-55.
- Blumenthal JA, Babyak MA, Wei J, et al. Usefulness of psychosocial treatment of mental stress-induced myocardial ischemia in men. *Am J Cardiol* 2002;89:164-68
- Blumenthal JA, Emery CF, Rejeski WJ. The effects of exercise training on psychosocial functioning after myocardial infarction. *J Cardiopulm Rehabil Prev* 1988b;8(5):183-93
- Blumenthal JA, Jiang W, Babyak MA, et al. Stress management and exercise training in cardiac patients with myocardial ischemia: effect on prognosis and evaluation of mechanisms. *Arch Int Med* 1997;157:2213-23
- Blumenthal JA, Rejeski WJ, Walsh-Riddle M, et al. Comparison of high- and low-intensity exercise training early after acute myocardial infarction. *Am J Cardiol*. 1988a;61(1):26-30
- Bohachick PA. Progressive relaxation training in cardiac rehabilitation: effect on psychologic variables. *Nurs Res* 1984;33:283-7

- Bolman C. Smoking cessation among patients hospitalized with cardiac disease: evaluation of a minimal-contact intervention (PhD-thesis). Maastricht: Universitaire Pers 2001
- Bonde JP, Munch-Hansen T, Agerbo E, et al. Job strain and ischemic heart disease: a prospective study using a new approach for exposure assessment. *J Occup Environ Med.* 2009;51(6):732-38.
- Bonde E, Breikss A, Hartford M. Effects of early rehabilitation on consumption of medical care during the first year after acute myocardial infarction in patients > or = 65 years of age. *Am J Cardiol* 1995;75(12):767-71
- Bostwick, J.M. and C.L. Sola, An updated review of implantable cardioverter/defibrillators, induced anxiety, and quality of life. *Psychiatr Clin North Am*, 2007. 30(4): 677-88.
- Bottcher M, Falk E. Pathology of the coronary arteries in smokers and non-smokers. *J Cardiovasc Risk* 1999;6:299-302
- Boyd KJ et al., Living with advanced heart failure: a prospective, community based study of patients and their carers. *Eur J Heart Fail* 2004;6(5): 585-91
- Braith RW, Welsch MA, Feigenbaum MS, et al. Neuroendocrine activation in heart failure is modified by endurance exercise training. *J Am Coll Cardiol* 1999;34:1170-5
- Brannstrom M, et al., Being a close relative of a person with severe, chronic heart failure in palliative advanced home care – a comfort but also a strain. *Scand J Caring Sci* 2007;21(3): 338-44
- Braunwald E. Heart Disease: A textbook of cardiovascular medicine (First edition). Lippincot Blackwell 1980
- Brechue W, Pollock MI. Exercise training for coronary artery disease in the elderly. *Clin Geriatr Med* 1996;12:207-29
- Briffa TG, Eckermann SD, Griffiths AD, et al. Cost-effectiveness of rehabilitation after an acute coronary event: a randomized controlled trial. *Med JAust* 2005;183(9):450-55
- Brink E, Brandstrom Y, Cliffordsson C, Herlitz J, Karlson BW. Illness consequences after myocardial infarction: problems with physical functioning and return to work. *J Adv Nurs* 2008;64(6):587-94
- Broadbent E, Ellis CJ, Thomas J, et al. Further development of an illness perception intervention for myocardial infarction patients: A randomized controlled trial. *J Psychosom Res* 2009;67:17-23
- Brostrom A et al. Congestive heart failure, spouses' support and the couple's sleep situation: a critical incident technique analysis. *J Clin Nurs* 2003;12(2):223-33-34.
- Brown N, Melville M, Gray D, et al. Quality of life four years after acute myocardial infarction: short form 36 scores compared with a normal population. *Heart* 1999;81(4):352-58
- Brügemann J, Hellemans IM. Commentaar: Reactie op de NVAB-richtlijn 'Handelen van de bedrijfsarts bij werknemers met ischemische hartziekten'. *TBV*, 15, 6, 2007.

- Bucher HC, Hengstler P, Schindler C, et al. N-3 polyunsaturated fatty acids in coronary heart disease: a meta-analysis of randomized controlled trials. *Am J Med* 2002;112:298-304
- Bundy C, Carroll D, Wallace L, et al. Psychological treatment of chronic stable angina pectoris. *Psychol & Health* 1994;10:69-77
- Bundy C, Carroll D, Wallace L, et al. Stress management and exercise training in chronic stable angina pectoris. *Psychol & Health* 1998;13:147-55
- Burg MM, et al. Low perceived social support and post-myocardial infarction prognosis in the enhancing recovery in coronary heart disease clinical trial: the effects of treatment. *Psychosom Med* 2005;67(6):879-88
- Burke JL, et al., The psychosocial impact of the implantable cardioverter defibrillator: a meta-analytic review. *Br J Health Psychol* 2003;8(Pt 2):165-78
- Burke LE, Dunbar-Jacob JM, Hill MN. Compliance with cardiovascular disease prevention strategies: a review of the research. *Ann Behav Med* 1997;19:239-63
- Burr ML, Fehily AM, Gilbert JF, et al. Effects of changes in fat, fish, and fibre intakes on death and myocardial reinfarction: diet and reinfarction trial (DART). *Lancet* 1989;2:757-61
- Callaerts-Vegh Z, Wenk M, Goebbels U, et al. Influence of intensive physical training on urinary nitrate elimination and plasma endothelin-1 levels in patients with congestive heart failure. *J Cardiopulm Rehabil* 1998;18:450-7
- Campbell N, Grimshaw J, Rawles J, et al. Cardiac rehabilitation: the agenda set by post myocardial infarction patients. *Health Educ J*;53(4):409-20
- Campbell N, Ritchie L, Rawles J, et al. Cardiac rehabilitation: the agenda set by post myocardial infarction patients. *Health Educ J* 1994;53:409-20
- Capomolla S, Febo O, Ceresa M, et al. Cost/utility in chronic heart failure: comparison between heart failure management program delivered by day-hospital and usual care. *J Am Coll Cardiol* 2002;40(7):1259-66
- Carinci F et al. Role of interactions between psychological and clinical factors in determining 6-month mortality among patients with acute myocardial infarction. Application of recursive partitioning techniques to the GISSI-2 database. Gruppo Italiano per lo Studio della Sopravvivenza nell' Infarto Miocardico. *Eur Heart J* 1997;18(5): 835-45
- Carney RM et al. Depression and late mortality after myocardial infarction in the Enhancing Recovery in Coronary Heart Disease (ENRICHD) study. *Psychosom Med* 2004;66(4): 466-74
- Carney RM et al. History of depression and survival after acute myocardial infarction. *Psychosom Med* 2009 ;71(3):253-59
- Carney RM, Freedland KE, Steinmeyer B et al. Depression and five year survival following acute myocardial infarction: a prospective study. *J Affect Disord* 2008;109:133-38

- Carson P, Phillips R, Lloyd M, et al. Exercise after myocardial infarction: a controlled trial. *J R Coll Physicians Lond* 1982;16(3):147-51
- CBO (Kwaliteits Instituut van de Geneeskunde CBO). Multidisciplinaire richtlijn cardiovasculair risicomangement 2006. Utrecht : Kwaliteitsinstituut voor de Gezondheidszorg CBO 2006.
- CBO (Kwaliteits Instituut van de Geneeskunde CBO). Richtlijn behandeling van tabaksverslaving 2003. Utrecht: Kwaliteitsinstituut voor de Gezondheidszorg CBO 2003.
- CBO (Kwaliteitsinstituut voor de Gezondheidszorg CBO). Evidence-based Richtlijnontwikkeling, Handleiding voor werkgroepleden. Kwaliteitsinstituut voor de Gezondheidszorg CBO 2007.
- CBS (Centraal Bureau voor de statistiek. Permanent onderzoek leefsituatie, gezondheid en arbeid 2003. Heerlen: CBS 2003 ([www.rivm.nl](http://www.rivm.nl))
- CBS. Bevolkingstrends, statistisch kwartaalblad over de demografie van Nederland, jaargang 51, 2<sup>e</sup> kartaal 2003. Heerlen/Voorburg: Centraal Bureau voor de Statistiek 2003
- Chandola T, Britton A, Brunner E, et al. Work stress and coronary heart disease: what are the mechanisms? *Eur Heart J* 2008;29(5):640-8
- Chandola T, Siegrist J, Marmot M. Do changes in effort-reward imbalance at work contribute to an explanation of the social gradient in angina? *Occ Environ Med* 2005;62(4):223-30
- Cheek F, Schrader G, Banham D, et al. Identification, course, and treatment of depression after admission for a cardiac condition: rationale and patient characteristics for the Identifying Depression As a Comorbid Condition (IDACC) project. *Am Heart J* 2003;146(6):978-84
- Chobanian AV, Bakris GL, Black HR, et al. The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: the JNC 7 report. *JAMA* 2003;289:2560-72
- Christensen KB, et al. The impact of psychosocial work environment factors on the risk of disability pension in Denmark. *Eur J Public Health* 2008;18(3):235-7
- Christensen KB, Feveile H, Labriola M, et al. The impact of psychosocial work environment factors on the risk of disability pension in Denmark. *Eur J Public Health*. 2008;18(3):235-7
- Chung MC et al. Posttraumatic stress and co-morbidity following myocardial infarction among older patients: the role of coping. *Aging Ment Health* 2008;12(1):124-33
- Clarke R, Frost C, Collins R, et al. Dietary lipids and blood cholesterol: quantitative meta-analysis of metabolic ward studies. *BMJ* 1997;314:112-7
- Clarke SP, Frasure-Smith N, Lespérance F, et al. Psychosocial factors as predictors of functional status at 1 year in patients with left ventricular dysfunction. *Research in Nursing & Health* 2000;23(4):290-300.
- Cleophas TJ. Wine, beer and spirits and the risk of myocardial infarction: a systematic review. *Biomed Pharmacother* 1999;53:417-23

- Cluitmans JJM, Vonken H, Bär F. Hartrevalidatie Deel 5: Hartrevalidatie en specialistische revalidatie. *Cardiologie* 1996;3:296-8
- Cluitmans JJM, Vonken H, Bär F. Hartrevalidatie Deel 6: Hartrevalidatie in revalidatiecentrum 'Hoensbroeck'. *Cardiologie* 1996;3:367-9
- Coats AJ, Adamopoulos S, Meyer TE, et al. Effects of physical training in chronic heart failure. *Lancet* 1990;335:63-6
- Coats AJ, Adamopoulos S, Meyer TE, et al. Effects of physical training in chronic heart failure. *Lancet* 1990;335:63-6
- Coats AJ, Adamopoulos S, Radaelli A, et al. Controlled trial of physical training in chronic heart failure. Exercise performance, hemodynamics, ventilation, and autonomic function. *Circulation* 1992;85:2119-31
- Coats AJ, Adamopoulos S. Neurohormonal mechanisms and the role of angiotensin-converting enzyme (ACE) inhibitors in heart failure. *Cardiovasc Drugs Ther* 1994;8(5):685-92
- Coats AJ, Adamopoulos S, Meyer TE, et al. Effects of physical training in chronic heart failure. *Lancet* 1990;335(8681):63-6
- Cobb LA, Weaver WD. Exercise: a risk for sudden death in patients with coronary heart disease. *J Am Coll Cardiol* 1986;7:215-9
- Cobelli F, Tavazzi L. Relative role of ambulatory and residential rehabilitation. *J Cardiovasc Risk* 1996;3:172-5
- Coleman MT, Newton KS. Supporting self-management in patients with chronic illness. *Am Fam Physician* 2005;72(8):1503-10
- Collins JA, Rice VH. Effects of relaxation intervention in phase II cardiac rehabilitation: replication and extension. *Heart & Lung* 1997;26:31-44
- Commissie Cardiovasculaire Preventie en Hartrevalidatie. Concept NVVC Praktijkrichtlijn Hartrevalidatie. Utrecht: NVVC, 2010
- Cooper AF, et al. Factors associated with cardiac rehabilitation attendance: a systematic review of the literature. *Clin Rehabil* 2002 ;16(5):541-52
- Cooper AF, Jackson G, Weinman J, et al. Factors associated with cardiac rehabilitation attendance: a systematic review of the literature. *Clin Rehabil* 2002;16(5):541-52
- Corrao G, Rubbiati L, Bagnardi V, et al. Alcohol and coronary heart disease: a meta-analysis. *Addiction* 2000;95:1505-23
- Cossette S, Fransure-Smith N., Lespérance F. Clinical implications of a reduction in psychological distress on cardiac prognosis in patients participating in a psychosocial intervention program. *Psychosom Med* 2001;63(2):257-66
- Cossette S, Fransure-Smith N, and Lespérance F. Clinical implications of a reduction in psychological distress on cardiac prognosis in patients participating in a psychosocial intervention program. *Psychosom Med* 2001;63(2):257-66



- Coster JE, Brügemann J, Oosterwijk MH, et al. Effectiviteit van klinische hartrevalidatie. *Hart Bulletin* 2002;33:3-6
- Cotugna N, Subar AF, Heimendinger J, et al. Nutrition and cancer prevention knowledge, beliefs, attitudes, and practices: the 1987 National Health Interview Survey. *J Am Diet Assoc* 1992;92:963-8
- Cowan MJ et al. Predictors of treatment response for depression and inadequate social support—the ENRICHD randomized clinical trial. *Psychother Psychosom* 2008;77(1): 27-37.
- Cowan MJ, Pike KC, Budzynski HK. Psychosocial nursing therapy following sudden cardiac arrest: impact on two-year survival. *Nurs Res* 2001;50:68-76
- Coyne JC, et al., Prognostic importance of marital quality for survival of congestive heart failure. *Am J Cardiol* 2001;88(5):526-29
- Criqui MH. Do known cardiovascular risk factors mediate the effect of alcohol on cardiovascular disease? *Novartis Found Symp* 1998;216:159-67
- Critchley JA, Capewell S. Mortality risk reduction associated with smoking cessation in patients with coronary heart disease: as systematic review. *JAMA* 2003;290:86-97
- Cupples ME, McKnight A. Randomised controlled trial of health promotion in general practice for patients at high cardiovascular risk. *BMJ* 1994;309:993-6
- Cupples S, et al. Report of the Psychosocial Outcomes Workgroup of the Nursing and Social Sciences Council of the International Society for Heart and Lung Transplantation: present status of research on psychosocial outcomes in cardiothoracic transplantation: review and recommendations for the field. *J Heart Lung Transplant* 2006;25(6):716-25.
- Curry SJ, Kristal AR, Bowen DJ. An application of the stage model of behavior change to dietary fat reduction. *Health Educ Res* 1992;7:97-105
- Cutler JA, Follmann D, Allender PS. Randomized trials of sodium reduction: an overview. *Am J Clin Nutr* 1997;65:643S-51S
- Dattilo AM, Kris-Etherton PM. Effects of weight reduction on blood lipids and lipoproteins: a meta-analysis. *Am J Clin Nutr* 1992;56:320-8
- Davidson KW et al. Enhanced depression care for patients with acute coronary syndrome and persistent depressive symptoms. *Arch Int Med* 2010;170(7):600-8
- Davies SJ, et al. Treatment of anxiety and depressive disorders in patients with cardiovascular disease. *BMJ* 2004;328(7445): 939-43
- de Backer G, Ambrosioni E, Borch-Johnsen K, et al. European guidelines on cardiovascular disease prevention in clinical practice: Third Joint Task Force of European and other Societies on Cardiovascular Disease Prevention in Clinical Practice (constituted by representatives of eight societies and by invited experts). *Eur Heart J* 2003;24:1601-10
- De Bacquer D, Pelfrene E, Clays E, et al. Perceived job stress and incidence of coronary events: 3-year follow-up of the Belgian Job Stress Project cohort. *Am J Epidemiol* 2005;161(5):434-41
- de Bono DP. Models of cardiac rehabilitation. *BMJ* 1998;316(7141):1329–1330

- De Groot NMS, et al. , The impact of an implantable cardioverter defibrillator: the Leiden follow-up study of ICD patients and their partners. *Neth J Cardiol* 2003;11:154-8.
- de Jong MJ, Chung ML, Roser LP, et al. A five-country comparison of anxiety early after acute myocardial infarction. *Eur J Cardiovasc Nurs*;3(2):129-34
- de Jonge P, et al. Only incident depressive episodes after myocardial infarction are associated with new cardiovascular events. *J Am Coll Cardiol* 2006;48(11): 2204-8.
- de Jonge P, et al. Symptom dimensions of depression following myocardial infarction and their relationship with somatic health status and cardiovascular prognosis. *Am J Psychiatry* 2006;163(1):138-44
- de Jonge P, Honig A, van Melle JP, et al. Nonresponse to treatment for depression following myocardial infarction: association with subsequent cardiac events. *Am J Psychiatry* 2007;164(9):1371-8
- de Loor SL, de Greef MHG, Jansma FFI, et al. Wat hartpatiënten beweegt. Een inventariserend onderzoek naar het aanbod van bewegingsactiviteiten voor hartpatiënten in Nederland. Bunnik/Groningen: 2001
- De Lorgeril M, Renaud S, Mamelle N, et al. TI-Mediterranean alpha-linolenic acid-rich diet in secondary prevention of coronary heart disease. *Lancet* 1994;343:1454-9
- De Lorgeril M, Salen P, Martin JL, et al. Mediterranean diet, traditional risk factors, and the rate of cardiovascular complications after myocardial infarction: final report of the Lyon Diet Heart Study. *Circulation* 1999;99:779-85
- De Lorgeril M, Salen P, Martin JL, et al. Wine drinking and risks of cardiovascular complications after recent acute myocardial infarction. *Circulation* 2002;106:1465-9
- De Rijk A, van Lierop B, Jansen N, Nijhuis F. Geen kwestie van motivatie maar van situatie. Een onderzoek naar man/vrouw verschillen in werkhervattin gedurende het eerste jaar na een ziekmelding. Doetinchem / 's Gravenhage: Elseviers bedrijfsinformatie / Ministerie van Sociale Zaken en Werkgelegenheid 2002
- De Rijk A, van Raak A, van der Made J. A new theoretical model for cooperation in public health settings: The RDIC-model. *Qual Health Res* 2007;17(8):1103-16
- De Rijk AE, et al. Gender differences in return to work patterns among sickness absentees and their associations with health: a prospective cohort study in The Netherlands. *Int J Rehabil Res*, 2008;31(4):327-36
- De Rijk AE. Verslag pilot Arbeid in het kader van PAAHR. Maastricht: Interne rapportage vakgroep Sociale Geneeskunde Universiteit Maastricht 2010
- DeBusk RF, Blomqvist CG, Kouchoukos NT, et al. Identification and treatment of low-risk patients after acute myocardial infarction and coronary-artery bypass graft surgery. *N Engl J Med* 1986;314(3):161-6
- DeBusk RF, Haskell WL, Miller NH, et al. Medically directed at-home rehabilitation soon after clinically uncomplicated acute myocardial infarction: a new model for patient care. *Am J Cardiol* 1985;55(4):251-7

- DeBusk RF, Houston N, Haskell W, et al. Exercise training soon after myocardial infarction. *Am J Cardiol* 1979;44(7):1223-9
- DeBusk RF, Miller NH, Superko HR, et al. A case-management system for coronary risk factor modification after acute myocardial infarction. *Ann Intern Med* 1994;120(9):721-9
- Delagardelle C, Feiereisen P, Autier P, et al. Strength/endurance training versus endurance training in congestive heart failure. *Med Sci Sports Exerc* 2002;34:1868-72
- Denke MA. Lack of efficacy of low-dose sitostanol therapy as an adjunct to a cholesterol-lowering diet in men with moderate hypercholesterolemia. *Am J Clin Nutr* 1995;61:392-6
- Denollet J, Brutsaert DL. Personality, disease severity, and the risk of long-term cardiac events in patients with a decreased ejection fraction after myocardial infarction. *Circulation* 1998;97(2): 167-73
- Denollet J, et al. Personality as independent predictor of long-term mortality in patients with coronary heart disease. *Lancet* 1996;347(8999):417-21
- Denollet J, et al. Unfavorable outcome of heart transplantation in recipients with type D personality. *J Heart Lung Transplant* 2007;26(2):152-8
- Denollet J, et al. Usefulness of type D personality in predicting five-year cardiac events above and beyond concurrent symptoms of stress in patients with coronary heart disease. *Am J Cardiol* 2006;97(7): 970-3
- Dew MA, DiMartini AF. Psychological disorders and distress after adult cardiothoracic transplantation. *J Cardiovasc Nurs* 2005;20(5 Suppl):S51-S66
- Dew MA, et al. Early post-transplant medical compliance and mental health predict physical morbidity and mortality one to three years after heart transplantation. *J Heart Lung Transplant* 1999;18(6):549-62
- Dickens C, et al. New onset depression following myocardial infarction predicts cardiac mortality. *Psychosom Med* 2008;70(4):450-5
- Dickens C, McGowan L, Percival C, et al. Depression is a risk factor for mortality after myocardial infarction: fact or artifact? *J Am Coll Cardiol* 2007;49(18):1834-40
- Dickens CM, et al., Lack of a close confidant, but not depression, predicts further cardiac events after myocardial infarction. *Heart* 2004;90(5):518-22
- DiMatteo MR, Lepper HS, Croghan TW. Depression is a risk factor for noncompliance with medical treatment: meta-analysis of the effects of anxiety and depression on patient adherence. *Arch Intern Med* 2000;160(14):2101-7
- DiMatteo, MR. Social support and patient adherence to medical treatment: a meta-analysis. *Health Psychol* 2004;23(2):207-18
- Dobbels F, De Geest S, Vanhees L, et al. Depression and the heart: a systematic overview of definition, measurement, consequences and treatment of depression in cardiovascular disease. *Eur J Cardiovasc Nurs* 2002;1(1):45-55

- Doerfler LA, JA Paraskos. Anxiety, posttraumatic stress disorder, and depression in patients with coronary heart disease: a practical review for cardiac rehabilitation professionals. *J Cardiopulm Rehabil* 2004;24(6):414-21
- Doerfler LA, Paraskos JA, Piniarski L. Relationship of quality of life and perceived control with posttraumatic stress disorder symptoms 3 to 6 months after myocardial infarction. *J Cardiopulm Rehabil* 2005b;25(3):166-72
- Doerfler LA, Paraskos JA. Post-traumatic stress disorder in patients with coronary artery disease: screening and management implications. *Can J Cardiol* 2005;21(8):689-97
- Doerfler LA, Paraskos JA. Post-traumatic stress disorder in patients with coronary artery disease: screening and management implications. *Can J Cardiol* 2005a;21(8):689-97.
- Doppegieter K. Privacywetgeving en omgaan met patiëntengegevens. Utrecht: KNMG Artsenfederatie Opleiding en registratie Domus medica 2002
- Dougherty CM, Pyper GP, Benoliel JQ. Domains of concern of intimate partners of sudden cardiac arrest survivors after ICD implantation. *J Cardiovasc Nurs* 2004;19(1): 21-31
- Dowlati Y et al. Efficacy and tolerability of antidepressants for treatment of depression in coronary artery disease: a meta-analysis. *Can J Psychiatry* 2010;55:91-9
- Dracup K, et al. Emotional well-being in spouses of patients with advanced heart failure. *Heart Lung* 2004;33(6):354-61
- Dressendorfer RH, Franklin BA, Cameron JL, et al. Exercise training frequency in early post-infarction cardiac rehabilitation. Influence on aerobic conditioning. *J Cardiopulm Rehabil* 1995;15(4):269-76
- Drexler H, et al. Alterations of skeletal muscle in chronic heart failure. *Circulation* 1992;85(5):1751-59
- Drexler H, et al. Endothelial function in congestive heart failure. *Am Heart J* 1993; 126(3 Pt 2):761-4
- Dubach P, Myers J, Dziekan G, et al. Effect of exercise training on myocardial remodelling in patients with reduced left ventricular function after myocardial infarction: application of magnetic resonance imaging. *Circulation* 1997;95:2060-7
- Dubach P, Myers J, Dziekan G, et al. Effect of high intensity exercise training on central hemodynamic responses to exercise in men with reduced left ventricular function. *J Am Coll Cardiol* 1997;29:1591-8
- Dubach P, Myers J, Dziekan G, et al. Effect of residential cardiac rehabilitation following bypass surgery. *Chest* 1995;108:1434-9
- Dugmore L, Tipson RJ, Philips MH, et al. Changes in cardiorespiratory fitness, psychological wellbeing, quality of life, and vocational status following a 12 month cardiac exercise rehabilitation programme. *Heart* 1999;81:359-66
- Dumont S, et al. Rehabilitation and the socio-occupational reintegration of workers who have had a myocardial infarct: a pilot study. *Can J Cardiol* 1999;15(4):453-61

- Dunbar-Jacob J. Models for changing patient behavior. *Am J Nurs* 2007;107(6 Suppl): 20-5
- Dusseldorp E, van Elderen T, Maes S, et al. A meta-analysis of psychoeducational programs for coronary heart disease patients. *Health Psychol* 1999;18(5):506-19
- Dziekan G, Myers J, Goebbels U, et al. Effects of exercise training on limb blood flow in patients with reduced ventricular function. *Am Heart J* 1998;136:22-30
- Eaker ED, Sullivan LM, Kelly-Hayes M, et al. Does job strain increase the risk for coronary heart disease or death in men and women? The Framingham Offspring Study. *Am J Epidemiol* 2004;159(10):950-8
- Earle A, Ayanian JZ, Heymann J. Work resumption after newly diagnosed coronary heart disease: findings on the importance of paid leave. *J Womens Health* 2006;15(4):430-41
- Eiber J, Mathes P. Partial inpatient cardiologic rehabilitation in an urban satellite center of a rehabilitation clinic: the Munich model. *Herz* 1999;24:63-6
- Eller NH, Netterstrom B, Gyntelberg F, et al. Work-related psychosocial factors and the development of ischemic heart disease: a systematic review. *Cardiol Rev* 2009;17(2):83-97
- Engblom E, Hietanen EK, Hamalainen H, et al. Exercise habits and physical performance during comprehensive rehabilitation after coronary artery bypass surgery. *Eur Heart J* 1992;13:1053-9
- Engblom E, Korpilahti K, Hamalainen H, et al. Quality of life and return to work 5 years after coronary artery bypass surgery. Long-term results of cardiac rehabilitation. *J Cardiopulm Rehabil* 1997;17:29-36
- EUROASPIRE I and II Study Group. Clinical reality of coronary prevention guidelines: a comparison of EUROASPIRE I and II in nine countries. *European Action on Secondary Prevention by Intervention to Reduce Events. Lancet* 2001;357(9261):995-1001
- EUROASPIRE II Study Group. Lifestyle and risk factor management and use of drug therapies in coronary patients from 15 countries; principal results from EUROASPIRE II Euro Heart Survey Programme. *Eur Heart J* 2001;22(7):554-72
- EUROASPIRE Study Group. A European Society of Cardiology survey of secondary prevention of coronary heart disease: principal results. *European Action on Secondary Prevention through Intervention to Reduce Events. Eur Heart J* 1997;18(10):1569-82
- Falger PR, Schouten EG. Exhaustion, psychological stressors in the work environment, and acute myocardial infarction in adult men. *J Psychosom Res.* 1992 Dec;36(8):777-86.
- Feigenbaum E, Carter E. Cardiac rehabilitation services Health technology assessment report, 1987, no 6. DHHS publication no. PHS 88-3427, Rockville, MD: National Center for Health Services Research and Health Care Technology Assessment, Public Health Service, U.S. Department of Health and Human Services 1987
- Ferrie JE, Shipley MJ, Davey Smith G, et al. Change in health inequalities among British civil servants: the Whitehall II study. *J Epidemiol Community Health* 2002;56(12):922-6

- Ferrie JE, Shipley MJ, Marmot MG, et al. Health effects of anticipation of job change and non-employment: longitudinal data from the Whitehall II study. *BMJ* 1995;11;311(7015):1264-9
- Ferrie JE, Shipley MJ, Marmot MG, Stansfeld SA, Smith GD. An uncertain future: the health effects of threats to employment security in white-collar men and women. *Am J Public Health* 1998;88(7):1030-6
- Fiore MC, Bailey WC, Cohen SJ, et al. Treating tobacco use and dependence, Clinical practice guideline. Rockville, MD: US Department of Health and Human Services. Public Health Service 2000
- Fleet R, Lavoie K, Beitman BD. Is panic disorder associated with coronary artery disease? A critical review of the literature. *J Psychosom Res* 2000;48(4-5):347-56
- Fletcher BJ, Dunbar SB, Felner JM, et al. Exercise testing and training in physically disabled men with clinical evidence of coronary artery disease. *Am J Cardiol* 1994;73(2):170-4
- Fletcher GF, Balady G, Blair SN, et al. Statement on exercise: benefits and recommendations for physical activity programs for all Americans. A statement for health professionals by the Committee on Exercise and Cardiac Rehabilitation of the Council on Clinical Cardiology, American Heart Association. *Circulation* 1996;94:857-62
- Fletcher GF, Balady G, Froelicher VF, et al. Exercise standards: a statement for healthcare professionals from the American Heart Association Writing Group. *Circulation* 1995;91:580-615
- Foster C, Pollock ML, Anholm JD, et al. Work capacity and left ventricular function during rehabilitation after myocardial revascularization surgery. *Circulation* 1984;69(4):748-55
- Foster C, Pollock MI, Anholm JD, et al. Work capacity and left ventricular function during rehabilitation after myocardial revascularization surgery. *Circulation* 1984;69:748-55
- Foster E et al. Task force 2: special health care needs of adults with congenital heart disease. *J Am Coll Cardiol* 200;37(5): 1176-83
- Franciosa JA, Park M, Levine, TB. Lack of correlation between exercise capacity and indexes of resting left ventricular performance in heart failure. *Am J Cardiol* 1981;7(1): 33-9.
- Franklin TL, Kolasa KM, Griffin K, et al. Adherence to very-low-fat diet by a group of cardiac rehabilitation patients in the rural southeastern United States. *Arch Fam Med* 1995;4:551-4
- Frasere-Smith N, Lespérance F, Gravel G, et al. Depression and health care costs during the first year following myocardial infarction. *J Psychosom Res* 2000;48(4-5):471-8
- Frasere-Smith N, Lespérance F, Prince RH, et al. Randomised trial of home-based psychosocial nursing intervention for patients recovering from myocardial infarction. *Lancet* 1997;350(9076):473-9

- Frasure-Smith N, Lespérance F, Talajic M. Depression and 18-month prognosis after myocardial infarction. *Circulation* 1995;91(4):999-1005
- Frasure-Smith N, Lespérance F, Talajic M. Depression following myocardial infarction. Impact on 6-month survival. *JAMA* 1993;270(15):1819-25
- Frasure-Smith N, Lespérance F. Reflections on depression as a cardiac risk factor. *Psychosom Med* 2005;67(suppl 1):S19-S25
- Frasure-Smith, N., et al., Long-term survival differences among low-anxious, high-anxious and repressive copers enrolled in the Montreal heart attack readjustment trial. *Psychosom Med* 2002;64(4): 571-9
- Frederiksen PM, Effects of physical training in children and adolescents with congenital heart disease. *Cardiol Young* 2000;10: 107-14
- Fredriksson M, Herlitz J, Nichol G. Variation in outcome in studies of out-of-hospital cardiac arrest: a review of studies conforming to the Utstein guidelines. *Am J Emerg Med* 2003;21(4): 276-81
- Freedland KE, Carney RM, Lustman PJ, et al. Major depression in coronary artery disease patients with vs. without a prior history of depression. *Psychosom Med* 1992;54(4):416-21
- Freedland KE, Skala JA, Carney RM, et al. Treatment of depression after coronary artery bypass surgery. A randomized controlled trial. *Arch Gen Psychiatry* 2009;66(4):387-396
- Friedmann E et al. Relationship of depression, anxiety, and social isolation to chronic heart failure outpatient mortality. *Am Heart J*, 2006. 152(5): 940 e1-8.
- Froelicher V, et al. A randomized trial of the effects of exercise training after CABG. *Arch Int Med* 1985;145:689-92
- Froelicher V, Jensen D, Genter F, et al. A randomized trial of exercise training in patients with coronary heart disease. *JAMA* 1984;252(10):1291-97
- Froom P, Cohen C, Rashcupkin J, et al. Referral to occupational medicine clinics and resumption of employment after myocardial infarction. *J Occup Environ Med* 1999;41(11):943-47
- Gagliardi JA, Prado NG, Marino JC, et al. Exercise training and heparin pretreatment in patient with coronary artery disease. *Am Heart J* 1996;132(5):946-51
- Gallagher JEJ, Hopkinson CA, Bennett P, et al. Effect of stress management on angina. *Psychol & Health* 1997;12:523-32
- Gallo LC et al. Marital status, marital quality, and atherosclerotic burden in postmenopausal women. *Psychosom Med* 2003;65(6):952-62
- Gallo WT, Teng HM, Falba TA, et al. The impact of late career job loss on myocardial infarction and stroke: a 10 year follow up using the health and retirement survey. *Occup Environ Med* 2006;63(10):683-7
- Gary RA. Combined exercise and CBT improves outcomes in patients with heart failure. *J Psychosom Res* 2010;69:119-131

- Geleijnse JM, Grobbee DE. Voeding en gezondheid - hypertensie. *Ned Tijdschr Geneesk* 2003;147:996-1000
- Gezondheidsraad. Overgewicht en obesitas (2003/07). 's Gravenhage; Gezondheidsraad 2003
- Gezondheidsraad. Voedingsnormen: energie, eiwitten, vetten en verteerbare koolhydraten (2001/19). 's Gravenhage; Gezondheidsraad 2001
- Giannattasio C, et al. Radial artery flow-mediated dilatation in heart failure patients: effects of pharmacological and nonpharmacological treatment. *Hypertension* 2001;38(6): 1451-5.
- Giannuzzi P, Tavazzi L, Temporelli PL, et al. Long-term physical training and left ventricular remodeling after anterior myocardial infarction: results of the Exercise in Anterior Myocardial Infarction (EAMI) trial. *J Am Coll Cardiol* 1993;22(7):1821-9
- Giannuzzi P, Temporelli PL, Tavazzi L, et al. EAMI-exercise training in anterior myocardial infarction: an ongoing multicenter randomized study. Preliminary results on left ventricular function and remodeling. *Chest* 1992;101(5 Suppl):315S-321S
- Gianuzzi P, Tavazzi L, Meyer K, et al. Recommendations for exercise training in chronic heart failure patients. *Eur Heart J*, 2001;22(2):125-35
- Gillis JG. Physicians' responsibilities on spaceship earth. *CMAJ* 1990;142(1):16
- Ginzburg K, Solomon Z, Bleich A. Repressive coping style, acute stress disorder, and posttraumatic stress disorder after myocardial infarction. *Psychosom Med* 2002;64(5):748-57
- Ginzburg K, Solomon Z, Bleich A. Repressive coping style, acute stress disorder, and posttraumatic stress disorder after myocardial infarction. *Psychosom Med*, 2002. 64(5): 748-57.
- Glassman AH, O'Connor CM, Califf RM, et al. Sertraline treatment of major depression in patients with acute MI or unstable angina. *JAMA* 2002;288(6):701-9
- Glazer KM et al. Psychological predictors of adherence and outcomes among patients in cardiac rehabilitation. *J Cardiopulm Rehabil* 2002;22(1): 40-6
- Glazer KM, Emery CF, Frid DJ, et al. Psychological predictors of adherence and outcomes among patients in cardiac rehabilitation. *J Cardiopulm Rehabil* 2002;22(1):40-6
- Goble AJ, Hare DL, Macdonald PS, et al. Effect of early programmes of high and low intensity exercise on physical performance after transmural acute myocardial infarction. *Br Heart J* 1991; 65(3): 126-31
- Goldberg RB, Mellies MJ, Sacks FM, et al. Cardiovascular events and their reduction with pravastatin in diabetic and glucose-intolerant myocardial infarction survivors with average cholesterol levels: subgroup analyses in the cholesterol and recurrent events (CARE) trial. The Care Investigators. *Circulation* 1998;98:2513-9
- Goldbert RB, Mellies MJ, Sacks FM, et al. Cardiovascular events and their reduction with pravastatin in diabetic and glucose-intolerant myocardial infarction survivors with



- average cholesterol levels: subgroup analyses in the chlesterol and recurrent events (CARE) trail. *Circulation* 1998;98:2513-19.
- Goldston K, Baillie AJ. Depression and coronary heart disease: a review of the epidemiological evidence, explanatory mechanisms and management approaches. *Clin Psychol Rev* 2008;28(2):288-306
- Gordon A, Tyni-Lenne R, Persson H, et al. Markedly improved skeletal muscle function with local muscle training in patients with chronic heart failure. *Clin Cardiol* 1996;19:568-74
- Gorgels AP, et al. Out-of-hospital cardiac arrest—the relevance of heart failure. The Maastricht Circulatory Arrest Registry. *Eur Heart J* 2003;24(13):1204-9
- Gortner SR. Nursing's syntax revisited: a critique of philosophies said to influence nursing theories. *Int J Nurs Stud* 1993;30(6):477-88
- Gottlieb SS, et al. Effects of exercise training on peak performance and quality of life in congestive heart failure patients. *J Card Fail* 1999;5(3):188-94
- Gottlieb SS, Fisher ML, Freudenberger R, et al. Effects of exercise training on peak performance and quality of life in congestive heart failure patients. *J Card Fail* 1999;5:188-94
- Goud R, de Keizer NF, ter Riet G, et al. Effect of guideline based computerised decision support on decision making of multidisciplinary teams: cluster randomised trial in cardiac rehabilitation. *BMJ* 2009;388:b144
- Grace SL, Abbey SE, Irvine J, et al. Prospective examination of anxiety persistence and its relationship to cardiac symptoms and recurrent cardiac events. *Psychother Psychosom* 2004;73(6):344-52
- Grace SL, Abbey SE, Pinto R, et al. Longitudinal course of depressive symptomatology after a cardiac event: effects of gender and cardiac rehabilitation. *Psychosom Med* 2005;67(1):52-8
- Grace SL, Abbey SE, Shnek ZM, et al. Cardiac rehabilitation I: review of psychosocial factors. *Gen Hosp Psychiatry*. 2002;24(3):121-6
- Grace SL, et al. Effect of depression on five-year mortality after an acute coronary syndrome. *Am J Cardiol* 2005;96(9):1179-85
- Grace SL, Leung YW, Stewart DE. A prospective examination of antidepressant use and its correlates in patients with acute coronary syndrome. *Psychosomatics* 2008;49(3):199-207
- Grady KL, et al., Team management of patients with heart failure: A statement for healthcare professionals from The Cardiovascular Nursing Council of the American Heart Association. *Circulation* 2000;102(19):2443-56
- Graham TP Jr., et al. 26<sup>th</sup> Bethesda conference: recommendations for determining eligibility for competition in athletes with cardiovascular abnormalities. Task Force 1: congenital heart disease. *J Am Coll Cardiol* 1994;24(4): 867-73

- Graudal NA, Galloe AM, Garred P. Effects of sodium restriction on blood pressure, renin, aldosterone, catecholamines, cholesterol, and triglycerides: a meta-analysis. *J Am Med Assoc* 1998;279:1383-91
- Green LA, Dickinson, P, Nease DE et al. Post-myocardial infarction depression clinical practice guideline panel. AAFP Guideline for the detection and management of post-myocardial depression. *Ann Fam Med* 2009;7:71-9.
- Greenland P, et al. Efficacy of cardiac rehabilitation services. With emphasis on patients after myocardial infarction. *Ann of Int Med* 1988;109:650-63
- Griffiths K, Camic PM, Hutton JM. Participant experiences of a mindfulness-based cognitive therapy group for cardiac rehabilitation. *J Health Psychol* 2009;14:675-681
- Grodzinski E, Jette M, Blumchen G, et al. Effects of a four-week training program on left ventricular function as assessed by radionuclide ventriculography. *J Cardiopulm Rehabil* 1987;7:518-24
- Haffner SM, Alexander CM, Cook TJ, et al. Reduced coronary events in simvastatin-treated patients with coronary heart disease and diabetes or impaired fasting glucose level: subgroup analyses in the Scandinavian Simvastatin Survival Study. *Arch Intern Med* 1999;159:2661-67.
- Hall JP, Wiseman VL, King MT, et al. Economic evaluation of a randomized trial of early return to normal activities versus cardiac rehabilitation after acute myocardial infarction. *Heart Lung Circ* 2002;11(1):10-8
- Hämäläinen H, Kallio V, Knuts L-R, et al. Community approach in rehabilitation and secondary prevention after acute myocardial infarction: results of a randomized clinical trial. *J Cardiopulm Rehabil Prev* 1991;11(4):221-26
- Hambrecht R, et al. Effect of exercise on coronary endothelial function in patients with coronary artery disease. *N Engl J Med* 2000;342(7):454-60
- Hambrecht R, et al. Regular physical exercise corrects endothelial dysfunction and improves exercise capacity in patients with chronic heart failure. *Circulation* 1998;98(24):2709-15
- Hambrecht R, Fiehn E, Weigl C, et al. Regular physical exercise corrects endothelial dysfunction and improves exercise capacity in patients with chronic heart failure. *Circulation* 1998;98:2709-15
- Hambrecht R, Fiehn E, Yu J, et al. Effects of endurance training on mitochondrial ultrastructure and fiber type distribution in skeletal muscle of patients with stable chronic heart failure. *J Am Coll Cardiol* 1997;29:1067-73
- Hambrecht R, Gielen S, Linke A, et al. Effects of exercise training on left ventricular function and peripheral resistance in patients with chronic heart failure: A randomized trial. *JAMA* 2000;283:3095-101
- Hambrecht R, Hilbrich L, Erbs S, et al. Correction of endothelial dysfunction in chronic heart failure: additional effects of exercise training and oral L-arginine supplementation. *J Am Coll Cardiol* 2000;35:706-13

- Hambrecht R, Niebauer J, Fiehn E, et al. Physical training in patients with stable chronic heart failure: effects on cardiorespiratory fitness and ultrastructural abnormalities of leg muscles. *J Am Coll Cardiol* 1995;25:1239-49
- Hambrecht R, Niebauer J, Marburger C, et al. Various intensities of leisure time physical activity in patients with coronary artery disease: effects on cardiorespiratory fitness and progression of coronary atherosclerotic lesions. *J Am Coll Cardiol* 1993;22:468-477
- Hanssen TA, et al. Anxiety and depression after acute myocardial infarction: an 18-month follow-up study with repeated measures and comparison with a reference population. *Eur J Cardiovasc Prev Rehabil* 2009;
- Hanssen TA, et al. Does a telephone follow-up intervention for patients discharged with acute myocardial infarction have long-term effects on health-related quality of life? A randomised controlled trial. *Journal of Clinical Nursing* 2009;18(9):1334-1345
- Hanssen TA, et al. Improving outcomes after myocardial infarction: a randomized controlled trial evaluating effects of a telephone follow-up intervention. *Eur J Cardiovasc Prev Rehabil* 2007;14(3):429-37
- Harrington D, et al. Skeletal muscle function and its relation to exercise tolerance in chronic heart failure. *J Am Coll Cardiol* 1997;30(7): 1758-64
- Hartley LH. Exercise and cardiac rehabilitation. *Proc N Engl Cardiovasc Soc* 1976;28:37-40
- Hase S, Douglas A. Effects of relaxation training on recovery from myocardial infarction. *Austr J Adv Nurses* 1987;5:18-27
- Haskell WL, Alderman EL, Fair JM, et al. Effects of intensive multiple risk factor reduction on coronary atherosclerosis and clinical cardiac events in men and woman with coronary artery disease. The Stanford Coronary Risk Intervention Project (SCRIP). *Circulation* 1994;89(3):975-90
- Haskell WL. Cardiovascular complications during exercise training of cardiac patients. *Circulation* 1978;57:920-4
- Haussler B, Keck M. Improvement in occupational rehabilitation of myocardial infarct patients--results of a model study in Rhineland-Pfalz. *Rehabilitation (Stuttg)* 1997;36(2):106-10
- Havranek EP et al. Predictors of the onset of depressive symptoms in patients with heart failure. *J Am Coll Cardiol*, 2004. 44(12): 2333-8.
- Haworth JE, *Eur J Heart Fail* 2005;7:803-808
- He FJ, MacGregor GA. Beneficial effects of potassium. *BMJ* 2001;323:497-501
- He J, Whelton PK, Appel LJ, et al. Long-term effects of weight loss and dietary sodium reductin on incidence of hypertension. *Hypertension* 2000;35:544-49.
- Hemingway H, Marmot M. Clinical Evidence: Psychosocial factors in the etiology and prognosis of coronary heart disease: systematic review of prospective cohort studies. *West J Med* 1999;171(5-6):342-350

- Hemingway H, Marmot M. Evidence based cardiology: psychosocial factors in the aetiology and prognosis of coronary heart disease. Systematic review of prospective cohort studies. *BMJ* 1999;318(7196): 1460-67
- Hemingway H, Vahtera J, Virtanen M, et al. Outcome of stable angina in a working population: the burden of sickness absence. *Eur J Cardiovasc Prev Rehabil* 2007;14(3):373-9
- Higgins HC, Hayes RL, McKenna KT. Rehabilitation outcomes following percutaneous coronary interventions (PCI). *Patient Educ Couns* 2001;43(3):219-30
- Hildebrand VH, Ooijendijk WTM, Stiggelbout M. Trendrapport Bewegen en Gezondheid 2000-2001. Hoofddorp: TNO Arbeid, TNO PG 2002
- Hildingh C, Fridlund B. A 3-year follow-up of participation in peer support groups after a cardiac event. *Eur J Cardiovasc Nurs* 2004;3(4):315-20
- Hillers TK, et al. Quality of life after myocardial infarction. *J Clin Epidemiol* 1994;47(11): 1287-96
- Hirschfeld RM, et al. The National Depressive and Manic-Depressive Association consensus statement on the undertreatment of depression. *JAMA* 1997; 277(4):333-40
- Hjermann I, Holme I, Leren P. Oslo Study Diet and Antismoking Trial. Results after 102 months. *Am J Med* 1986;80:7-11
- Hjermann I, Velve Byre K, Holme I, et al. Effect of diet and smoking intervention on the incidence of coronary heart disease. *Lancet* 1981;2:1303-10
- Hofgren C, et al. Two years after cardiac arrest; cognitive status, ADL function and living situation. *Brain Inj* 2008;22(12):972-8
- Hofman-Bang C, Lisspers J, Nordlander R, et al. Two-year results of a controlled study of residential rehabilitation for patients treated with percutaneous transluminal coronary angioplasty. A randomized study of a multifactorial programme. *Eur Heart J* 1999;20(20):1465-74
- Hogan BE, Linden W, Najarian B. Social support interventions: do they work? *Clin Psychol Rev* 2002;22:381-440.
- Honig A et al. Treatment of post-myocardial infarction depressive disorder: a randomized, placebo-controlled trial with mirtazapine. *Psychosom Med* 2007;69(7):606-13
- Honig A. Depression following a heart infarct and increased risk of death. *Ned Tijdschr Geneesk* 2000;144(27):1307-10
- Honig A. et al. Depression following a first heart infarct; similarities with and differences from 'ordinary' depression. *Ned Tijdschr Geneesk*, 1997. 141(4): 196-99
- Hooper L, Summerbell CD, Higgins JP, et al. Dietary fat intake and prevention of cardiovascular disease: systematic review. *BMJ* 2001;322:757-63
- Horlick L, Cameron R, Firor W, et al. The effects of education and group discussion in the post-myocardial infarction patient. *J Psychosom Res* 1984;28(6):485-92

- Horlick L, Cameron R, Firor W, et al. The effects of education and group discussion in the post-myocardial infarction patient. *J Psychosom Res* 1984;28:485-92
- Hornig B, Maier V, Drexler H. Physical training improves endothelial function in patients with chronic heart failure. *Circulation*, 1996; 3(2):210-4
- Hossack KF, et al. Cardiac arrest associated with supervised cardiac rehabilitation. *J Cardiac Rehabil* 1982;2:402-8
- Hu FB, Stampfer MJ, Manson JE, et al. Dietary intake of alpha-linolenic acid and risk of fatal ischemic heart disease among women. *Am J Clin Nutr* 1999;69:890-7
- Huffman JC, Celano CM, Januzzi. Review. The relationship between depression, anxiety, and cardiovascular outcomes in patients with acute coronary syndromes. *Neuropsychiatric Disease and Treatment* 2010;6:123-36
- Huffman JC, Smith FA, Blais MA, et al. Recognition and treatment of depression and anxiety in patients with acute myocardial infarction. *Am J Cardiol* 2006;98(3):319-24
- Huisman M, Van Lenthe F, Avendano M, et al. The contribution of job characteristics to socioeconomic inequalities in incidence of myocardial infarction. *Soc Sci Med*. 2008;66(11):2240-52
- Hung J, Gordon EP, Houston N, et al. Changes in rest and exercise myocardial perfusion and left ventricular function 3 to 26 weeks after clinically uncomplicated acute myocardial infarction: effects of exercise training. *Am J Cardiol* 1984;54(8):943-50
- Jalowiec A, Grady KL, White-Williams C. Functional status one year after heart transplant. *J Cardiopulm Rehabil Prev* 2007; 27(1):24-32; 33-34
- Janeway D. An integrated approach to the diagnosis and treatment of anxiety within the practice of cardiology. *Cardiol Rev* 2009;17(1): 36-43
- Janssen N, van den Heuvel WP, Beurskens AJ, et al. The Demand-Control-Support model as a predictor of return to work. *Int J Rehabil Res* 2003;26(1):1-9
- Johnston M, et al. Impact on patients and partners of inpatient and extended cardiac counseling and rehabilitation: a controlled trial. *Psychosom Med* 1999;61(2):225-33
- Jolliffe JA, Rees K, Taylor RS, et al. Exercise-based rehabilitation for coronary heart disease. *Cochrane Database Syst Rev* 2001;1:CD001800
- Jones DA, West RR. Psychological rehabilitation after myocardial infarction: multicentre randomised controlled trial. *BMJ* 1996;313(7071):1517-21
- Joshi KJ, Hu FB, Manson JE, et al. The effect of fruit and vegetable intake on risk for coronary heart disease. *Ann Intern Med* 2001;134:1106-14
- Kamphuis HC, et al. A 12-month quality of life assessment of cardiac arrest survivors treated with or without an implantable cardioverter defibrillator. *Europace* 2002;4(4):417-25
- Kamphuis M et al. Health related quality of life and health status in adult survivors with previously operated complex congenital heart disease. *Heart* 2002;87(4):356-62

- Kamphuis M, Vogels T, Ottenkamp J, et al. Employment in adults with congenital heart disease. *Arch Pediatr Adolesc Med* 2002;156(11):1143-8
- Kaptein KI, de Jonge P, van den Brink RH, et al. Course of depressive symptoms after myocardial infarction and cardiac prognosis: a latent class analysis. *Psychosom Med* 2006;68(5):662-8
- Karlsson MR et al. Effects of expanded cardiac rehabilitation on psychosocial status in coronary artery disease with focus on type D characteristics. *J Behav Med* 2007;30(3):53-61
- Karner AM, Dahlgren MA, Bergdahl B. Rehabilitation after coronary heart disease: spouses' views of support. *J Adv Nurs* 2004;46(2):204-11
- Kasper EK, Gerstenblith G, Heftner G, et al. A randomized trial of the efficacy of multidisciplinary care in heart failure outpatients at high risk of hospital readmission. *J Am Coll Cardiol* 2002;39(3):471-80
- Kavanagh T, et al. Cardiorespiratory responses to exercise training after orthotopic cardiac transplantation. *Circulation* 1988;77(1):162-71
- Kavanagh T, et al. Long-term cardiorespiratory results of exercise training following cardiac transplantation. *Am J Cardiol* 2003;91(2):190-4
- Kavanagh T, Shephard RJ, Pandit V, et al. Exercise and hypnotherapy in the rehabilitation of the coronary patient. *Arch Phys Med Rehab* 1970;Oct:578-87
- Kavanagh T, YM, Campbell R, et al., Marathon running after cardiac transplantation: a case report. *J Cardiopulmon Rehabil* 1986;6:16-20
- Kemper HCG, Ooijendijk WTM, Stiggelbout M, et al. Consensus over de Nederlandse Norm Gezond Bewegen. *TSG* 2000;78:180-83
- Kemper HCG, Ooijendijk WTM, Stiggelbout M, et al. Consensus over de Nederlandse Norm Gezond Bewegen. *TSG* 2000;78:180-3
- Keteyian SJ, Brawner CA, Schairer JR, et al. Effects of exercise training on chronotropic incompetence in patients with heart failure. *Am Heart J* 1999;138:233-40
- Keteyian SJ, Levine AB, Brawner CA, et al. Exercise training in patients with heart failure. A randomized, controlled trial. *Ann Intern Med* 1996;124:1051-7
- Kettunen S, et al. Myocardial infarction, spouses' reactions and their need of support. *J Adv Nurs* 1999;30(2):479-88
- Kiilavuori K, Naveri H, Leinonen H, et al. The effect of physical training on hormonal status and exertional hormonal response in patients with chronic congestive heart failure. *Eur Heart J* 1999;20:456-64
- Kiilavuori K, Naveri H, Salmi T, et al. The effect of physical training on skeletal muscle in patients with chronic heart failure. *Eur J Heart Fail* 2000;2:53-63
- Kiilavuori K, Sovijarvi A, Naveri H, et al. Effect of physical training on exercise capacity and gas exchange in patients with chronic heart failure. *Chest* 1996;110:985-91

- Kimble LP. Impact of cardiac symptoms on self-reported household task performance in women with coronary artery disease. *J Cardiopulm Rehabil* 2001;21(1):18-23
- King CN, Senn MD. Exercise testing and prescription. Practical recommendation for the sedentary. *Sports Med* 1996;5:326-36
- Kitchiner D. Physical activities in patients with congenital heart disease. *Heart*, 1996. 76(1):6-7
- Kivimaki M, Vahtera J, Elovainio M, et al. Human costs of organizational downsizing: comparing health trends between leavers and stayers. *Am J Community Psychol* 2003; 32(1-2):57-67
- Kivimaki M, Virtanen M, Elovainio M, et al. Work stress in the etiology of coronary heart disease—a meta-analysis. *Scand J Work Environ Health* 2006;32(6):431-42
- Klempt Hw, Roman C, Ylinen T, et al. Partial inpatient cardiologic after-care (article in German). *Herz* 1999;24:32-7
- Ko DT, et al. Beta-blocker therapy and symptoms of depression, fatigue, and sexual dysfunction. *JAMA* 2002;288(3):351-7
- Kobashigawa JA, et al. A controlled trial of exercise rehabilitation after heart transplantation. *N Engl J Med* 1999;340(4): 272-7
- Koertge J, et al. Effects of a stress management program on vital exhaustion and depression in women with coronary heart disease: a randomized controlled intervention study. *J Intern Med* 2008;263(3):281-93
- Komajda M, et al. The EuroHeart Failure Survey programme—a survey on the quality of care among patients with heart failure in Europe. Part 2: treatment. *Eur Heart J* 2003;24(5):464-74
- Kop WJ, et al. Vital exhaustion predicts new cardiac events after successful coronary angioplasty. *Psychosom Med* 1994;56(4):281-7
- Kornitzer M, deSmet P, Sans S, et al. Job stress and major coronary events: results from the Job Stress, Absenteeism and Coronary Heart Disease in Europe study. *Eur J Cardiovasc Prev Rehabil* 2006;13(5):695-704
- Kovacs AH, et al. Adults with congenital heart disease: psychological needs and treatment preferences. *Congenit Heart Dis* 2009;4(3):139-46
- Kovacs AH, et al. Depression and anxiety in adult congenital heart disease: predictors and prevalence. *Int J Cardiol* 2009;137(2):158-64
- Krachler M, Lindschinger M, Eber B, et al. Trace elements in coronary heart disease: impact of intensified lifestyle modification. *Biol Trace Elem Res* 1997;60(3):175-85
- Kraus WE, Houmard JA, Duscha BD, et al. Effects of the amount and intensity of exercise on plasma lipoproteins. *N Engl J Med* 2002;347(19):1483-92
- Kris-Etherton PM, Harris WS, Appel LJ. Fish consumption, fish oil, omega-3 fatty acids, and cardiovascular disease. *Circulation* 2002;106:2747-57

- Kris-Etherton PM, Hecker KD, et al. Bioactive compounds in foods: their role in the prevention of cardiovascular disease and cancer. *Am J Med* 2002;113:71S-88S
- Krishnan KR, et al. Comorbidity of depression with other medical diseases in the elderly. *Biol Psychiatry* 2002;52(6):559-88
- Kromhout D, Menotti A, Kesteloot H, et al. Prevention of coronary heart disease by diet and lifestyle: evidence from prospective cross-cultural, cohort, and intervention studies. *Circulation* 2002;105:893-98.
- Kromhout D, Nauta ILD. Preventie van coronaire hartziekten door voeding en leefstijl. *Klinische Cardiologie* 2003;1:5-9
- Krumholz HM, Amatrudda J, Smith GL, et al. Randomized trial of an education and support intervention to prevent readmission of patients with heart failure. *J Am Coll Cardiol* 2002;39(1):83-89
- Krumholz HM, et al. Prognostic importance of emotional support for elderly patients hospitalized with heart failure. *Circulation* 1998;97(10):958-64
- Kruse M, Hochstrasser S, Zwisler AD, et al. Comprehensive cardiac rehabilitation: a cost assessment based on a randomized clinical trial. *Int J Technol Assess Health Care* 2006;22(4):478-83
- Kulcu DG, et al. The effect of cardiac rehabilitation on quality of life, anxiety and depression in patients with congestive heart failure. A randomized controlled trial, short-term results. *Eura Medicophys* 2007;43(4):489-97
- Kuper H, Adami H, Theorell T, et al. Psychosocial determinants of coronary heart disease in middle-aged women: a prospective study in Sweden. *Am J Epidemiol* 2006;164(4):349-57
- Kutzleb J, Reiner D. The impact of nurse-directed patient education on quality of life and functional capacity in people with heart failure. *J Am Acad Nurse Pract* 2006;18(3):116-23.
- Ladwig KH, Kieser M, König J, et al. Affective disorders and survival after acute myocardial infarction. Results from the post-infarction late potential study. *Eur Heart J* 1991;12(9):959-64
- LaFontaine T. The role of lipid management by diet and exercise in the progression, stabilization, and regression of coronary artery atherosclerosis. *J Cardiopulm Rehabil* 1995;15(4):262-8
- Lallukka T, Martikainen P, Reunanen A, et al. Associations between working conditions and angina pectoris symptoms among employed women. *Psychosom Med* 2006;8(2):348-54
- Lancaster T, Silagy C, Fowler G. Training health professionals in smoking cessation (Cochrane Review). In: *The Cochrane Library*, Issue 3, 2002. Oxford: Update Software
- Lane D, Carroll D, Ring C, et al. Do depression and anxiety predict recurrent coronary events 12 months after myocardial infarction? *QJM* 2000;93(11):739-44



- Lane D, Carroll D, Ring C, et al. Mortality and quality of life 12 months after myocardial infarction: effects of depression and anxiety. *Psychosom Med* 2001;63(2):221-30
- Lane D, Carroll D, Ring C, et al. The prevalence and persistence of depression and anxiety following myocardial infarction. *Br J Health Psychol* 2002;7(Pt 1):11-21
- Lavie CJ, Milani RV, Littman AB. Benefits of cardiac rehabilitation and exercise training in secondary coronary prevention in the elderly. *J Am Coll Cardiol* 1993;22(3):678-83
- Lavie CJ, Milani RV. Adverse psychological and coronary risk profiles in young patients with coronary artery disease and benefits of formal cardiac rehabilitation. *Arch Intern Med* 2006;166(17):1878-83.
- Lavie CJ, Milani RV. Benefits of Cardiac Rehabilitation and Exercise Training in Older Persons. *Am J Geriatr Cardiol* 1995b;4(4):42-48
- Lavie CJ, Milani RV. Benefits of cardiac rehabilitation and exercise training in elderly women. *Am J Cardiol* 1997;79(5):664-6
- Lavie CJ, Milani RV. Cardiac rehabilitation and depression. *Am J Cardiol* 2004;93(8):1080
- Lavie CJ, Milani RV. Effects of cardiac rehabilitation and exercise training programs on coronary patients with high levels of hostility. *Mayo Clin Proc* 1999;74(10):959-66
- Lavie CJ, Milani RV. Effects of cardiac rehabilitation programs on exercise capacity, coronary risk factors, behavioral characteristics, and quality of life in a large elderly cohort. *Am J Cardiol* 1995a;76(3):177-9
- Lavie CJ, Milani RV. Prevalence of hostility in young coronary artery disease patients and effects of cardiac rehabilitation and exercise training. *Mayo Clin Proc* 2005;80(3):335-42
- Lavie CV, Milani RV. Impact of aging on hostility in coronary patients and effects of cardiac rehabilitation and exercise training in elderly persons. *Am J Geriatr Cardiol* 2004;13(3):125-30
- Law M. Plant sterol and stanol margarines and health. *BMJ* 2000;320:861-4
- Law MR, Morris JK. By how much does fruit and vegetable consumption reduce the risk of ischaemic heart disease: response to commentary. *Eur J Clin Nutr* 1999;53:903-4
- Law MR, Wald NJ, Thompson SG. By how much and how quickly does reduction in serum cholesterol concentration lower risk of ischaemic heart disease? *BMJ* 1994;308:367-72
- Law MR, Wald NJ. Risk factor thresholds: their existence under scrutiny. *BMJ* 2002;324:1570-6
- Ledwidge M, Barry M, Cahill J, et al. Is multidisciplinary care of heart failure cost-beneficial when combined with optimal medical care? *Eur J Heart Fail* 2003;5(3):381-9
- Lee AJ, Strickler GK, Shepard DS. The economics of cardiac rehabilitation and lifestyle modification: a review of literature. *J Cardiopulm Rehabil Prev* 2007;27(3):135-42

- Lee S, Colditz GA, Berkman LF, et al. Prospective study of job insecurity and coronary heart disease in US women. *Ann Epidemiol* 2004;14(1):24-30
- Lemaitre R, Siscovick D, Raghunathan T. Leisure-time physical activity and the risk of primary cardiac arrest. *Arch Intern Med* 1999;159:686-90
- Lemaitre RN, King IB, Mozaffarian D, et al. N-3 polyunsaturated fatty acids, fatal ischemic heart disease, and nonfatal myocardial infarction in older adults: the Cardiovascular Health Study. *Am J Clin Nutr* 2003;77:319-25
- Lemonick MD, Park A. Eat Smart. *Time* 1999 Jul 19: 44-6.
- Lespérance F, et al. Effects of citalopram and interpersonal psychotherapy on depression in patients with coronary artery disease: the Canadian Cardiac Randomized Evaluation of Antidepressant and Psychotherapy Efficacy (CREATE) trial. *JAMA* 2007;297(4):367-79.
- Lett HS, Blumenthal JA, Babyak MA, et al. In response to “perceived social support predicts outcomes following myocardial infarction: A call for screening?” *Health Psychol* 2008; 27(1):1-3
- Lett HS, et al. Social support and coronary heart disease: epidemiologic evidence and implications for treatment. *Psychosom Med* 2005;67(6): 869-78
- Lett HS, et al. Social support and prognosis in patients at increased psychosocial risk recovering from myocardial infarction. *Health Psychol* 2007;26(4): 418-27
- Lichtman JH, Bigger JT Jr, Blumenthal JA, et al. Depression and coronary heart disease: recommendations for screening, referral, and treatment: a science advisory from the American Heart Association Prevention Committee of the Council on Cardiovascular Nursing, Council on Clinical Cardiology, Council on Epidemiology and Prevention, and Interdisciplinary Council on Quality of Care and Outcomes Research: endorsed by the American Psychiatric Association. *Circulation* 2008;118(17):1768-75
- Lim C, et al. The neurological and cognitive sequelae of cardiac arrest. *Neurology* 2004;63(10):1774-78
- Linden W, Stossel C, Maurice J. Psychosocial interventions for patients with coronary artery disease: a meta-analysis. *Arch Intern Med* 1996;156(7):745-52
- Linden, W., M.J. Phillips, and J. Leclerc, Psychological treatment of cardiac patients: a meta-analysis. *Eur Heart J* 2007;28(24):2972-84.
- Linke A, et al. Endothelial dysfunction in patients with chronic heart failure: systemic effects of lower-limb exercise training. *J Am Coll Cardiol* 2001;37(2):392-97
- Linke A, Schoene N, Gielen S, et al. Endothelial dysfunction in patients with chronic heart failure: systemic effects of lower-limb exercise training. *J Am Coll Cardiol* 2001;37:392 -7
- Lip GY, et al. Psychological interventions for depression in adolescent and adult congenital heart disease. *Cochrane Database Syst Rev*, 2003(3): CD004394
- LIPID study group (Long-Term Intervention with Pravastatin in Ischaemic Diseases study group. *N Engl J Med* 1998;339:1349-57.

- Lipkin DP, et al. Abnormalities of skeletal muscle in patients with chronic heart failure. *Int J Cardiol* 1988;18(2):187-95
- Lowensteyn I, Coupal L, Zowall H, et al. The cost-effectiveness of exercise training for the primary and secondary prevention of cardiovascular disease. *J Cardiopulm Rehabil* 2000;20(3):147-55
- Lukkarinen H, Kyngas H. Experiences of the onset of coronary artery disease in a spouse. *Eur J Cardiovasc Nurs* 2003;2(3):189-94
- Lundgren-Nilsson A, et al. The first year after successful cardiac resuscitation: function, activity, participation and quality of life. *Resuscitation* 2005;66(3):285-89
- Luttik ML, et al. For better and for worse: Quality of life impaired in HF patients as well as in their partners. *Eur J Cardiovasc Nurs* 2005;4(1):11-14
- Luttik ML, et al. Marital status, quality of life, and clinical outcome in patients with heart failure. *Heart Lung* 2006;35(1):3-8
- Luttik ML, et al. Quality of life in partners of people with congestive heart failure: gender and involvement in care. *J Adv Nurs* 2009;65(7):1442-51
- Luttik ML, et al. The importance and impact of social support on outcomes in patients with heart failure: an overview of the literature. *J Cardiovasc Nurs* 2005;20(3):162-9
- Maeland J, Havik O. Return to work after a myocardial infarction: the influence of background factors, work characteristics and illness severity. *Scand J Soc Med* 1986;14:183-95
- Maeland JG, Havik OE. Psychological predictors for return to work after a myocardial infarction. *J Psychosom Res* 1987;31(4):471-81
- Maiorana A, O'Driscoll G, Dembo L, et al. Exercise training, vascular function, and functional capacity in middle-aged subjects. *Med Sci Sports Exerc* 2001;33:2022-8
- Malinauskienė V, Theorell T, Grazuleviciene R, et al. Low job control and myocardial infarction risk in the occupational categories of Kaunas men, Lithuania. *J Epidemiol Commun Health* 2004;58(2):131-5
- Mancini DM, et al. Contribution of intrinsic skeletal muscle changes to <sup>31</sup>P NMR skeletal muscle metabolic abnormalities in patients with chronic heart failure. *Circulation* 1989;80(5):1338-46
- Mark DB, Lam LC, Lee KL, et al. Identification of patients with coronary disease at high risk for loss of employment. A prospective validation study. *Circulation* 1992; 86(5):1485-94
- Maron BJ, Mitchell JH. Revised eligibility recommendations for competitive athletes with cardiovascular abnormalities. *J Am Coll Cardiol* 1994;24(4): 848-50
- Marra S, Paolillo V, Spadaccini F, et al. Long-term follow-up after a controlled randomized post-myocardial infarction rehabilitation programme: effects on morbidity and mortality. *Eur Heart J* 1985;6(8):656-63
- Martens EJ, et al. Cardiac history, prior depression and personality predict course of depressive symptoms after myocardial infarction. *Psychol Med* 2008;38(2):257-64

- Martensson J, Dracup K, Fridlund B. Decisive situations influencing spouses' support of patients with heart failure: a critical incident technique analysis. *Heart Lung* 2001;30(5): 341-50.
- Martin CR, Lewin RJ, Thompson DR. A confirmatory factor analysis of the Hospital Anxiety and Depression Scale in coronary care patients following acute myocardial infarction. *Psychiatry Res* 2003;120(1):85-94
- Mauro AM. Exploring uncertainty and psychosocial adjustment after cardioverter defibrillator implantation. *J Cardiovasc Nurs* 2008;23(6):527-35
- May GA, Nagle FJ. Changes in rate-pressure product with physical training of individuals with coronary artery disease. *Phys Ther* 1984;64(9):1361-6
- Mayou R, MacMahon D, Sleight P, et al. Early rehabilitation after myocardial infarction. *Lancet* 1981;2(8260-61):1399-402
- Mayou R. Rehabilitation after heart attack. *BMJ* 1996;313:1498-9
- Mayou RA, Gill D, Thompson DR, et al. Depression and anxiety as predictors of outcome after myocardial infarction. *Psychosom Med* 2000;62(2):212-9
- Mayou RA, Thompson DR, Clements A, et al. Guideline-based early rehabilitation after myocardial infarction. A pragmatic randomized controlled trial. *J Psychosom Res* 2002;52(2):89-95
- McCormick A, Fleming D, Charlton J. Morbidity statistics from general practice. Fourth national study 1991-92 . Office of Population Censuses and Surveys (Series MB5 No 3.). London: HMSO 1995.
- McCormick A, Fleming D, Charlton J. Morbidity statistics from general practice: fourth national study 1991 - 1992: a study carried out by the Royal College of General Practitioners, Office of Population Censuses and Surveys, and the Department of Health. London: HMSO 1995
- McKechnie R, Mosca L. Physical activity and coronary heart disease: prevention and effect on risk factors. *Cardiol Rev* 2003;11:21-5
- McKee, G. Are there meaningful longitudinal changes in health related quality of life-SF36, in cardiac rehabilitation patients? *European Journal of Cardiovascular Nursing* 2009;8(1): 40-47
- McKelvie RS, Teo KK, Roberts R, et al. Effects of exercise training in patients with heart failure: the Exercise Rehabilitation Trial (EXERT). *Am Heart J* 2002;144:23-30
- McLean S, Timmins F. An exploration of the information needs of spouse/partner following acute myocardial infarction using focus group methodology. *Nurs Crit Care* 2007;12(3):141-50
- McLeod A. Later management of documented ischaemic heart disease: secondary prevention and rehabilitation. *Brit Med Bull* 2001;59:113-33.
- Mead WF. Successful resuscitation of two near simultaneous cases of cardiac arrest with an overview of fifteen cases occurring during supervised exercise. *Circulation* 1976;53:187-9

- Meershoek A, Krumeich A, Desain L. Arbeidsongeschiktheid, reïntegratie en etniciteit. Amsterdam: Het Spinhuis 2005
- Meland E, Maeland JG, Laerum E. The importance of self-efficacy in cardiovascular risk factor change. *Scand J Public Health* 1999;27(1):11-7
- Mendes de Leon CF, et al., Psychosocial characteristics and recurrent events after percutaneous transluminal coronary angioplasty. *Am J Cardiol* 1996;77(4):252-55
- Menotti A, Blackburn H, Kromhout D, et al. cardiovascular risk factors as determinants of 25-year all-cause mortality in the seven countries study. *Eur J Epidemiol* 2001;17:337-46
- Mensink RP, Katan MB. Effect of dietary fatty acids on serum lipids and lipoproteins. A meta-analysis of 27 trials. *Arterioscler Thromb* 1992;12:911-9
- Meyer F, White E. Alcohol and nutrients in relation to colon cancer in middle-aged adults. *Am J Epidemiol* 1993;138:225-36
- Meyer K, Gornandt L, Schwaibold M, et al. Predictors of response to exercise training in severe chronic congestive heart failure. *Am J Cardiol* 1997;80:56-60
- Meyer K, Schwaibold M, Westbrook S, et al. Effects of short-term exercise training and activity restriction on functional capacity in patients with severe chronic congestive heart failure. *Am J Cardiol* 1996;78:1017-22
- Meyer TE, Casadei B, Coats AJ, et al. Angiotensin-converting enzyme inhibition and physical training in heart failure. *J Intern Med* 1991;230:407-13
- Meyer TE, Casadei B, Coats AJ, et al. Angiotensin-converting enzyme inhibition and physical training in heart failure. *J Intern Med* 1991;230(5):407-13
- Michalsen A, et al. Psychological and quality-of-life outcomes from a comprehensive stress reduction and lifestyle program in patients with coronary artery disease: results of a randomized trial. *Psychother Psychosom* 2005;74(6):344-52
- Midgley JP, Matthew AG, Greenwood CM, et al. Effect of reduced dietary sodium on blood pressure: a meta-analysis of randomized controlled trials. *JAMA* 1996;275:1590-7
- Miller ER, III, Appel LJ, Risby TH. Effect of dietary patterns on measures of lipid peroxidation: results from a randomized clinical trial. *Circulation* 1998;98:2390-5
- Miller NH, Haskell WL, Berra K, et al. Home versus group exercise training for increasing functional capacity after myocardial infarction. *Circulation* 1984;70(4):645-9
- Mirowski M. The automatic implantable cardioverter-defibrillator, an overview. *J Am Coll Cardiol* 1998;6(2):461-6.
- Mital A, et al. Accelerating the return to work (RTW) chances of coronary heart disease (CHD) patients: part 1--development and validation of a training programme. *Disabil Rehabil* 2000;22(13-14):604-20
- Mitchell JH, Haskell WL, Raven PB. Classification of sports. *J Am Coll Cardiol* 1994;24(4):864-6

Mittag O, et al. Outcomes of cardiac rehabilitation with versus without a follow-up intervention rendered by telephone (Luebeck follow-up trial): overall and gender-specific effects. *Int J Rehabil Res* 2006;29(4):295-302

Mittag O, Kolenda KD, Nordman KJ, et al. Return to work after myocardial infarction/coronary artery bypass grafting: patients' and physicians' initial viewpoints and outcome 12 months later. *Soc Sci Med* 2001;52(9):1441-50

Mittleman M, Maclure M, Tofler G. Triggering of acute myocardial infarction by heavy physical exertion. Protection against triggering by regular exertion. *N Engl J Med* 1993;329:1677-83

Moller J, Theorell T, de Faire U, et al. Work related stressful life events and the risk of myocardial infarction. Case-control and case-crossover analyses within the Stockholm heart epidemiology programme (SHEEP). *J Epidemiol Community Health* 2005; 59(1):23-30

Molloy GJ et al. Marital status and cardiac rehabilitation attendance: a meta-analysis. *Eur J Cardiovasc Prev Rehabil* 2008b;15(5): 557-61.

Molloy GJ et al. Practical support predicts medication adherence and attendance at cardiac rehabilitation following acute coronary syndrome. *J Psychosom Res* 2008a;65(6):581-6

Molloy GJ, et al. Social networks and partner stress as predictors of adherence to medication, rehabilitation attendance, and quality of life following acute coronary syndrome. *Health Psychol* 2008c;27(1):52-8

Molloy GJ, Johnston DW, Witham MD. Family caregiving and congestive heart failure. Review and analysis. *Eur J Heart Fail* 2005;7(4):592-603

Mols F, Martens EJ, Denollet J. Type D personality and depressive symptoms are independent predictors of impaired health status following acute myocardial infarction. *Heart* 2010;96(1): 30-5.

Mookadam F, Arthur HM, Social support and its relationship to morbidity and mortality after acute myocardial infarction: systematic overview. *Arch Intern Med* 2004; 64(14):1514-18.

Moser DK, Dracup K, Evangelista LS, et al. Comparison of prevalence of symptoms of depression, anxiety, and hostility in elderly patients with heart failure, myocardial infarction, and a coronary artery bypass graft. *Heart and Lung* 2010; 39(5):378-85

Moser DK, Dracup K. Role of spousal anxiety and depression in patients' psychosocial recovery after a cardiac event. *Psychosom Med*, 2004. 66(4): 527-32.

Moser DK, Riegel B, McKinley S, et al. Impact of anxiety and perceived control on in-hospital complications after acute myocardial infarction. *Psychosom Med* 2007;69(1):10-6

Moser DK, Watkins JF. Conceptualizing self-care in heart failure: a life course model of patient characteristics. *J Cardiovasc Nurs*, 2008. 23(3): 205-18; 219-20.

- Moser, D.K., et al., An international perspective on gender differences in anxiety early after acute myocardial infarction. *Psychosom Med*, 2003. 65(4): 511-6.
- Moser, DK, Dracup K. Is anxiety early after myocardial infarction associated with subsequent ischemic and arrhythmic events? *Psychosom Med* 1996;58(5):395-401
- Moss AJ, et al. Improved survival with an implanted defibrillator in patients with coronary disease at high risk for ventricular arrhythmia. Multicenter Automatic Defibrillator Implantation Trial Investigators. *N Engl J Med* 1996;335(26):1933-40
- Mosterd WL, Rosen PWF. Multidisciplinaire richtlijn Chronisch hartfalen. Netherlands Multidisciplinary guidelines CHF 2002
- Mozaffarian D, Kumanyika SA, Lemaitre RN, et al. Cereal, fruit, and vegetable fiber intake and the risk of cardiovascular disease in elderly individuals. *JAMA* 2003;289:1659-66
- Mullen PD, Mains DA, Velez R. A meta-analysis of controlled trials of cardiac patient education. *Patient Educ Couns* 1992;19:143-62
- Muller-Nordhorn J, Gehring J, Kulig M, et al. Return to work after cardiologic rehabilitation. *Soz Präventivmed*. 2003;48(6):370-8.
- Multidisciplinaire richtlijn Depressie. Eerste revisie. 2010.
- Muntwyler J, Hennekens CH, Buring JE, et al. Mortality and light to moderate alcohol consumption after myocardial infarction. *Lancet* 1998;352:1882-5
- Myers J, Gianrossi R, Schwitzer J, et al. Effect of exercise training on postexercise oxygen uptake kinetics in patients with reduced ventricular function. *Chest* 2001;120:1206-11
- Myers J, Wagner D, Schertler T, et al. Effects of exercise training on left ventricular volumes and function in patients with nonischemic cardiomyopathy: application of magnetic resonance myocardial tagging. *Am Heart J* 2002;144:719-25
- Mykletun A, Bjerkeset O, Dewey M, et al. Anxiety, depression, and cause-specific mortality: the HUNT study. *Psychosom Med* 2007;69(4):323-31
- Myrtek M, et al. Factors associated with work resumption: a 5 year follow-up with cardiac patients. *Int J Cardiol* 1997;59(3):291-97
- Naylor MD, Brooten DA, Campbell RL, et al. Transitional care of older adults hospitalized with heart failure: a randomized, controlled trial. *J Am Geriatr Soc* 2004;52(5):675-84
- NCEP-expert panel (National Cholesterol Education Program expert panel). Executive summary of the Third Report of on Detection, Evaluation, and Treatment of High Blood Cholesterol in Adults (Adult Treatment Panel III). *J Am Med Assoc* 2001; 285:2486-97.
- Nederlands WHO-FIC Collaborating Centre. Nederlandse vertaling van de 'International Classification of Functioning, Disability and Health'. Houten: RIVM National Institute for Public Health and the Environment 2002 (p. 291)
- Nederlandse Hartstichting. Hart en Vaatziekten in Nederland 2001. 's Gravenhage: Nederlandse Hartstichting 2001

Nederlandse Hartstichting. Hartrevalidatie: PEP-module. Handleiding voor professionals. (www.hartstichting.nl)

Nelson D, Baer PE, Cleveland SE, et al. Six-month follow-up of stress management training versus cardiac education during hospitalization for acute myocardial infarction. *J Cardiopulm Rehabil* 1994;14:384-90

Ness AR, Powles JW. Fruit and Vegetables, and Cardiovascular Disease: a review. *Int J Epidemiology* 1997;26:1-13

Newton M, Mutrie N, McCarthur JD. The effects of exercise in a coronary rehabilitation programme. *Scott Med J* 1991;36(2):38-41

NHLBI (National Heart and Lung and Blood Institute). Clinical Guidelines on the identification, evaluation, and treatment of overweight and obesity in adults. The evidence report 1998. Bethesda: National Institute of Health, National Heart and Lung and Blood Institute 1998

NHS (National Health Service) Centre for Reviews and Dissemination. Cardiac rehabilitation. *Eff Health Care* 1998; 4(4). (www.york.ac.uk)

Nicholson, A., H. Kuper, H. Hemingway. Depression as an aetiologic and prognostic factor in coronary heart disease: a meta-analysis of 6362 events among 146 538 participants in 54 observational studies. *Eur Heart J* 2006;27(23):2763-74

Nieboer AP, et al. Spousal caregivers' activity restriction and depression: a model for changes over time. *Soc Sci Med* 1998;47(9):1361-71

Nieminen H, et al. Long-term results of pediatric cardiac surgery in Finland: education, employment, marital status, and parenthood. *Pediatrics* 2003;112(6): 1345-50

Nieuwland W, Berkhuisen MA, van Veldhuisen DJ, et al. Differential effects of high-frequency versus low-frequency exercise training in rehabilitation of patients with coronary artery disease. *J Am Coll Cardiol* 2000;36(1):202-7

NIH (National Institute of Health) .Physical Activity and Cardiovascular Health. National Institute of Health Consensus Developmental Panel on Physical Activity and Cardiovascular Health. *JAMA* 1996;276:241-6

Nissen NK, Madsen M, Olsen AD et al. Health service interventions targeting relatives of heart patients: a review of the literature. *Scand J Public Health* 2008;36(8):818-26

Nolwajka AJ, Kostuk WJ, Rechnitzer PA, et al. Exercise and human collateralization: an angiographic and scintigraphic assessment. *Circulation* 1979;60(1):114-21

Nomura K, Nakao M, Sato M, et al. The association of the reporting of somatic symptoms with job stress and active coping among Japanese white-collar workers. *J Occup Health* 2007;49(5):370-5

NVD (Nederlandse Vereniging van Diëtisten). *Artsenwijzer diëtetiek* 2003. Houten: NVD 2003 (www.nvdietist.nl)

NVVC (Nederlandse Vereniging voor Cardiologie) werkgroep 'Congenitale cardiologie bij volwassenen'. *Volwassenen met congenitale hartziekten in Nederland: richtlijnen* 2000. 's Gravenhage: de Nederlandse Hartstichting 2000



- Nvvc, CBO, NHS, MISGZ. Consensus hart- en vaatziekten en sportbeoefening 1988.
- NVVG. Verzekeringsgeneeskundige protocollen COPD Chronisch hartfalen. Utrecht: NVVG 2007.
- NVVG. Verzekeringsgeneeskundige protocollen Whiplash / Aspecifieke lage rugpijn / Hartinfarct. Utrecht: NVVG 2008.
- O'Connell AM, Crawford MH, Abrams J. Heartfailure disease management in an indigent population. *Am Heart J* 2001;141(2):254-8
- O'Connor CM, Glassman AH, Harrison DJ. Pharmacoeconomic analysis of sertraline treatment of depression in patient with unstable angina or a recent myocardial infarction. *J Clin Psychiatry* 2005;66(3):346-52
- O'Connor GT, Buring JE, Yusuf S, et al. An overview of randomized trials of rehabilitation with exercise after myocardial infarction. *Circulation* 1989 Aug;80(2):234-44
- O'Neill C, Normand C, Cupples M, et al. A comparison of three measures of perceived distress: results from a study of angina patients in general practice in Northern Ireland. *J Epidemiol Community Health* 1996;50:202-6
- O'Reilly SM, Grubb N, O'Carroll REO. Long-term emotional consequences of in-hospital cardiac arrest and myocardial infarction. *Br J Clin Psychol* 2004;43:83-95
- O'Rourke Á, Hampson SE. Psychosocial outcomes after an MI: evaluation of two approaches to rehabilitation. *Psychol Health Med* 1999;4(4):393-402
- Oberman A, Fletcher GF, Lee J, et al. Efficacy of high-intensity exercise training of left ventricular ejection fraction in men with coronary artery disease (the Training Level Comparison Study). *Am J Cardiol* 1995;76(10):643-7
- Ohm D. Entspannungstraining und Hypnose bei Patienten mit koronaren Herzkrankheit in der stationären Rehabilitation. Regensburg: Roderer Verlag 1987
- Ohtsubo M, Yonezawa K, Nishijima H, et al. Metabolic abnormality of calf skeletal muscle is improved by localised muscle training without changes in blood flow in chronic heart failure. *Heart* 1997;78:437-43
- Oldridge N, Furlong W, Feeny D, et al. Economic evaluation of cardiac rehabilitation soon after acute myocardial infarction. *Am J Cardiol* 1993;72(7):154-61
- Oldridge N, Guyatt G, Jones N, et al. Effects on quality of life with comprehensive rehabilitation after acute myocardial infarction. *Am J Cardiol* 1991;67(13):1084-9
- Oldridge NB, Gottlieb M, Guyatt G, et al. Predictors of health related quality of life with cardiac rehabilitation after acute myocardial infarction. *J Cardiopulm Rehabil* 1998;18(2):95-103
- Oldridge NB, Guyatt GH, Fischer ME, et al. Cardiac rehabilitation after myocardial infarction. Combined experience of randomized clinical trials. *JAMA* 1988;260(7):945-50
- Oldridge NB, McCartney N, Hicks A, et al. Improvement in maximal isokinetic cycle ergometry with cardiac rehabilitation. *Med Sci Sports Exerc* 1989;21(3):308-12

- Oldridge NB. Cardiac rehabilitation and risk factor management after myocardial infarction: clinical and economic evaluation. *Wien Klin Wochenschr* 1997;109(S2):6-16
- Oldridge NB. Comprehensive cardiac rehabilitation: is it cost-effective? *Eur Heart J* 1998;19 SupplO:O42-50
- O'Neill C, Normand C, Cupples M, et al. A comparison of three measures of perceived distress: results from a study of angina patients in general practice in Northern Ireland. *J Epidemiol Community Health* 1996;50(2):202-6
- Ornish D, Brown SE, Scherwitz LW, et al. Lifestyle changes and heart disease. *Lancet* 1990;336:741-2
- Ornish D, Scherwitz LW, Billings JH, et al. Intensive lifestyle changes for reversal of coronary heart disease. *JAMA* 1998;280(23):2001-7
- Ornish D, Scherwitz LW, Doody RS, et al. Effects of stress management training and dietary changes in treating ischemic heart disease. *JAMA* 1983;249(1):54-9
- Orth-Gomér K et al. Stress reduction prolongs life in women with coronary disease: the Stockholm Women's Intervention Trial for Coronary Heart Disease (SWITCHD). *Circ Cardiovasc Qual Outcomes* 2009;2:25-32
- Orth-Gomér, K., et al., Marital stress worsens prognosis in women with coronary heart disease: the Stockholm Female Coronary Risk Study. *JAMA* 2000;284(23):3008-14
- Papadakis S, Oldridge NB, Coyle D, et al.. Economic evaluation of cardiac rehabilitation: a systematic review. *Eur J Cardiovasc Prev Rehabil* 2005;12(6):513-20
- Parashar S, Rumsfeld JS, Spertus JA, et al. Time course of depression and outcome of myocardial infarction. *Arch Intern Med.* 2006;166(18):2035-43
- Paris W, White-Williams C. Social adaptation after cardiothoracic transplantation: a review of the literature. *J Cardiovas Nurs* 2005;20(5S):S67-S73
- Parker GB, Owen CA, Brotchie HL, et al. The impact of differing anxiety disorders on outcome following an acute coronary syndrome: time to start worrying? *Depress Anxiety* 2010 Mar;27(3):302-9
- Parker, G.B., et al., The impact of differing anxiety disorders on outcome following an acute coronary syndrome: time to start worrying? *Depress Anxiety*, 2009. 27(3): 302-309.
- Pate RR, Pratt M, Blair SN, et al. Physical activity and health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* 1995;273(5):402-7
- Pate RR, Pratt M, Blair SN, et al. Physical activity and health: a recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* 1995;273:402-7
- Pate RR, Pratt M, Blair SN, et al. Physical activity and public health. A recommendation from the Centers for Disease Control and Prevention and the American College of Sports Medicine. *JAMA* 1995;273:402-7

- Pech-Amsellem MA, Myara I, Storogenko M, et al. Enhanced modifications of low-density lipoproteins (LDL) by endothelial cells from smokers: a possible mechanism of smoking-related atherosclerosis. *Cardiovasc Res* 1996;31:975-83
- Pedersen SS, et al. Clustering of poor device acceptance and Type D personality is associated with increased distress in Danish cardioverter-defibrillator patients. *Pacing Clin Electrophysiol* 2009d;32(1):29-36
- Pedersen SS, et al. Clustering of psychosocial risk factors enhances the risk of depressive symptoms 12-months post percutaneous coronary intervention. *Eur J Cardiovasc Prev Rehabil* 2008;15(2):203-9
- Pedersen SS, et al. Increased anxiety in partners of patients with a cardioverter-defibrillator: the role of indication for ICD therapy, shocks, and personality. *Pacing Clin Electrophysiol* 2009c;32(2):184-92
- Pedersen SS, et al. Poor perceived social support in implantable cardioverter defibrillator (ICD) patients and their partners: cross-validation of the multidimensional scale of perceived social support. *Psychosomatics* 2009b;50(5):461-67
- Pedersen SS, et al. Risk of chronic anxiety in implantable defibrillator patients: A multi-center study. *Int J Cardiol* 2009a;
- Pedersen SS, et al. Type D personality predicts death or myocardial infarction after bare metal stent or sirolimus-eluting stent implantation: a Rapamycin-Eluting Stent Evaluated at Rotterdam Cardiology Hospital (RESEARCH) registry substudy. *J Am Coll Cardiol* 2004;44(5):997-1001
- Pedersen SS, van den Broek KC, Sears SF Jr. Psychological intervention following implantation of an implantable defibrillator: a review and future recommendations. *Pacing Clin Electrophysiol* 2007;30(12):1546-54
- Pelle, A.J., et al., Psychological predictors of prognosis in chronic heart failure. *J Card Fail* 2008;14(4):341-50
- Penninx BWJH, van Dyck R. Stand van zaken: Depressie en somatische comorbiditeit. *Ned Tijdschr Geneesk* 2010; 154:A1784
- Penque S, Petersen B, Arom K, et al. Early discharge with home health care in the coronary artery bypass patient. *Dimens Crit Care Nurs* 1999;18(6):40-8
- Perk J, Alexanderson K. Swedish Council on Technology Assessment in Health Care (SBU). Chapter 8. Sick leave due to coronary artery disease or stroke. *Scand J Public Health* 2004;63S:181-206
- Peter R, Hammarstrom A, Hallqvist J, et al. Does occupational gender segregation influence the association of effort-reward imbalance with myocardial infarction in the SHEEP study? *Int J Behav Med* 2006;13(1):34-43
- Peter R, Siegrist J, Hallqvist J, et al. Psychosocial work environment and myocardial infarction: improving risk estimation by combining two complementary job stress models in the SHEEP Study. *J Epidemiol Community Health* 2002;56(4):294-300

- Petrie KJ, et al. Role of patients' view of their illness in predicting return to work and functioning after myocardial infarction: longitudinal study. *BMJ* 1996;312(7040):1191-94.
- Petrie KJ, Weinman J, Sharpe N, et al. Role of patients' view of their illness in predicting return to work and functioning after myocardial infarction: longitudinal study. *BMJ* 1996;312(7040):1191-4
- Pfund A, et al. Coronary intervention and occupational rehabilitation--a prospective, randomized intervention study. *Z Kardiol* 2001;0(9):655-60
- Phillips L, Harrison T, Houck P. Return to work and the person with heart failure. *Heart Lung* 2005;34(2):79-88
- Picard MH, et al. Cost-benefit analysis of early return to work after uncomplicated acute myocardial infarction. *Am J Cardiol* 1989;63(18):1308-14
- Piepoli M, et al. Contribution of muscle afferents to the hemodynamic, autonomic, and ventilatory responses to exercise in patients with chronic heart failure: effects of physical training. *Circulation* 1996;93(5):940-52
- Piepoli MF, Corrà U, Benzer W, et al. Secondary prevention through cardiac rehabilitation: from knowledge to implementation. A position paper from the Cardiac Rehabilitation Section of the European Association of Cardiovascular Prevention and Rehabilitation. *Eur J Cardiovasc Prev Rehabil* 2010;17(1):1-17
- Piepoli MF, Flather M, Coats AJ. Overview of studies of exercise training in chronic heart failure: the need for a prospective randomized multicentre European trial. *Eur Heart J* 1998;19(6):830-41
- Pignone MP, Ammerman A, Fernandez L, et al. Counseling to promote a healthy diet in adults: a summary of the evidence for the U.S. Preventive Services Task Force. *Am J Prev Med* 2003;24:75-92
- Pina IL, et al. Exercise and heart failure: A statement from the American Heart Association Committee on exercise, rehabilitation, and prevention. *Circulation* 2003;107(8):1210-25
- Poikolainen K. It can be bad for the heart, too-drinking patterns and coronary heart disease. *Addiction* 1998;93:1757-9
- Polackova J, Bockova E, Sedivec V. Autogenic training: application in secondary prevention of myocardial infarction. *Activ Nerv Sup* 1982;24-3:178-80
- Powell JT. Vascular damage from smoking: disease mechanisms at the arterial wall. *Vasc Med* 1998;3:21-28
- Prescott E, Hippe M, Schnohr P, et al. Smoking and risk of myocardial infarction in women and men: longitudinal population study. *BMJ* 1998;316:1043-47
- Prochaska JO, DiClemente CC. Stages and processes of self-change of smoking: toward an integrative model of change. *J Consult Clin Psychol* 1983;51:390-95

- Pu CT, Johnson MT, Forman DE, et al. Randomized trial of progressive resistance training to counteract the myopathy of chronic heart failure. *J Appl Physiol* 2001;90:2341-50
- Rees K, et al. Psychological interventions for coronary heart disease. *Cochrane Database Syst Rev* 2004;(2):CD002902
- Regan KL. Depression treatment with selective serotonin reuptake inhibitors for the postacute coronary syndrome population: a literature review. *J Cardiovasc Nursing* 2008;23:489-496
- Reincke C, Verstappen F. Biologische mechanismen ter verklaring van het gustige effect van lichamelijke inspanning op hart- en vaat ziekten. *Geneeskunde en Sport* 2001;34:110-18
- Remme WJ, Swedberg K. Guidelines for the diagnosis and treatment of chronic heart failure. *Eur Heart J* 2001;22(17):1527-60
- Rich MW, Beckham V, Wittenberg C, et al. A multidisciplinary intervention to prevent the readmission of elderly patients with congestive heartfailure. *N Engl J Med* 1995;333(18):1190-95
- Richtlijnen Hartrevalidatie deel I. Den Haag. Revalidatie Commissie. Nederlandse Vereniging Voor Cardiologie en de Nederlandse Hartstichting. 's Gravenhage: Nederlandse Hartstichting 1995
- Richtlijnen Hartrevalidatie deel II. Den Haag. Revalidatie Commissie Nederlandse Vereniging Voor Cardiologie en de Nederlandse Hartstichting. 's Gravenhage: Nederlandse Hartstichting 1995/1996
- Ridder DTD, Schreurs, KMG. Coping en sociale steun van chronisch zieken. Zoetermeer: NCCZ-reeks 1994
- Riegel B, Carlson B, Glaser D, et al. Which patients with heart failure respond best to multidisciplinary disease management? *J Card Fail* 2000;6(4):290-9
- Riegel B, Carlson B. Is individual peer support a promising intervention for persons with heart failure? *J Cardiovasc Nurs* 2004;19(3):174-83
- Riese H, Van Doornen LJ, Houtman IL, et al. Job strain in relation to ambulatory blood pressure, heart rate, and heart rate variability among female nurses. *Scand J Work Environ Health* 2004;30(6):477-85
- Rigotti NA, Munafo MR, Murphy MFG, et al. Interventions for smoking cessation in hospitalized patients (Cochrane Review). In: *The Cochrane Library*, Issue 1, 2003. Oxford: Update Software
- Rimm EB, Ascherio A, Giovannucci E, et al. Vegetable, fruit, and cereal fiber intake and risk of coronary heart disease among men. *JAMA* 1996;275:447-51
- Rimm EB, Williams P, Fosher K, et al. Moderate alcohol intake and lower risk of coronary heart disease: meta-analysis of effects on lipids and haemostatic factors. *BMJ* 1999;319:1523-8

- Robertson KA, Kayhko K. Cost analysis of an intensive home follow-up program for first-time post-myocardial infarction patients and their families. *J Can Ass Crit Care Nurses* 2001; 12(4): 25-31
- Robertson KA, Kayhko K. Cost analysis of an intensive home follow-up program for first-time post-myocardial infarction patients and their families. *Dynamics* 2001;12(4):25-31
- Roest AM, Martens EJ, Denollet J, et al. Prognostic association of anxiety post myocardial infarction with mortality and new cardiac events: a meta-analysis. *Psychosom Med* 2010;72(6):563-9
- Rohrbaugh MJ, et al. Couples coping with congestive heart failure: role and gender differences in psychological distress. *J Fam Psychol* 2002;16(1):3-13
- Rohrbaugh MJ, Shoham V, Coyne JC. Effect of marital quality on eight-year survival of patients with heart failure. *Am J Cardiol* 2006;98(8):1069-72
- Roine RO, Kajaste S, Kaste M. Neuropsychological sequelae of cardiac arrest. *JAMA*, 1993;269(2): 237-42
- Rothenbacher D, Hahmann H, Wüsten B, et al. Symptoms of anxiety and depression in patients with stable coronary heart disease: prognostic value and consideration of pathogenetic links. *Eur J Cardiovasc Prev Rehabil* 2007;14(4):547-54
- Rozanski A, Blumenthal JA, Davidson KW, et al. The epidemiology, pathophysiology, and management of psychosocial risk factors in cardiac practice: the emerging field of behavioral cardiology. *J Am Coll Cardiol* 2005;45(5):637-51
- Rozanski A, Blumenthal JA, Kaplan J. Impact of psychological factors on the pathogenesis of cardiovascular disease and implications for therapy. *Circulation* 1999; 99(16):2192-217
- Rozanski A, et al. The epidemiology, pathophysiology, and management of psychosocial risk factors in cardiac practice: the emerging field of behavioral cardiology. *J Am Coll Cardiol* 2005;45(5): 637-51
- Ruberman W, Weinblatt E, Goldberg JD, et al. Psychosocial influences on mortality after myocardial infarction. *New Engl J Med* 1984;311(9):552-9
- Rubins HB, Robins SJ, Collins D, et al. Gemfibozil for the secondary prevention of coronary heart disease in men with levels of high-density lipoprotein. *N Engl J Med* 1999;34:410-8
- Rugulies R. Depression as a predictor for coronary heart disease. A review and meta-analysis. *Am J Prev Med* 2002;23(1):51-61
- Rutledge T, Reis VA, Linke SE, Greenberg BH, Mills PJ. Depression in heart failure a meta-analytic review of prevalence, intervention effects, and associations with clinical outcomes. *J Am Coll Cardiol* 2006;48(8):1527-37
- Saab 2009] et al, The impact of cognitive behavioral group training on event-free survival in patients with myocardial infarction: The ENRICHD experience. *J Psychosom res* 2009;67:45-56.

- Sacks FM, Svetkey LP, Vollmer WM, et al. Effects on blood pressure of reduced dietary sodium and the Dietary Approaches to Stop Hypertension (DASH) diet. DASH-Sodium Collaborative Research Group. *N Engl J Med* 2001;344:3-10
- Salmoirago-Blotcher E, Ockene IS. Methodological limitations of psychosocial interventions in patients with an implantable cardioverter-defibrillator (ICD) A systematic review. *BMC Cardiovasc Disord* 2009;29;9:56
- Salonen JT. Stopping smoking and long-term mortality after acute myocardial infarction. *Br Heart J* 1980;43:463-9
- Samkange-Zeeb F, Altenhoner T, Berg G, et al. Predicting non-return to work in patients attending cardiac rehabilitation. *Int J Rehabil Res* 2006; 29(1):43-9
- Saris WH, Blair SN, van Baak MA, et al. How much physical activity is enough to prevent unhealthy weight gain? Outcome of the IASO 1st Stock Conference and consensus statement. *Obes Rev* 2003;4:101-14
- Sato I, Nishida M, Okita K, et al. Beneficial effect of stopping smoking on future cardiac events in male smokers with previous myocardial infarction. *Jpn Circ J* 1992;56:217-22
- Sato I, Nishida M, Okita K, et al. Beneficial effect of stopping smoking on future cardiac events in male smokers with previous myocardial infarction. *Jpn Circ J* 1992;56:217-22
- Saunders MM, Family caregivers need support with heart failure patients. *Holist Nurs Pract* 2003;17(3):136-42
- Sauve MJ, et al. Factors associated with cognitive recovery after cardiopulmonary resuscitation. *Am J Crit Care* 1996;5(2):127-39
- Sayers SL, et al. Social support and self-care of patients with heart failure. *Ann Behav Med* 2008;35(1):70-9
- Scherer M, et al. Disease-specific quality of life in primary care patients with heart failure. *Z Arztl Fortbild Qualitatssich* 2007a;101(3):185-90
- Scherer M, et al. Psychological distress in primary care patients with heart failure: a longitudinal study. *Br J Gen Pract* 2007b;57(543):801-7
- Scherer M, et al. Psychosocial determinants for frequent primary health care utilisation in patients with heart failure. *GMS Psychosoc Med* 2008;5:Doc02
- Schneiderman N, Saab PG et al. Psychosocial treatment within sex by ethnicity subgroups in the Enhancing Recovery in Coronary Heart Disease clinical trial. *Psychosom Med* 2004;66:475-483
- Schnohr P, Parner J, Lange P. Mortality in joggers: population based study of 4658 men. *BMJ* 2000;321:602-3
- Schofield DJ, Shrestha RN, Passey ME, et al. Chronic disease and labour force participation among older Australians. *Med J Aust* 2008;189(8):447-50
- Scholz U, et al. Physical activity and depressive symptoms in cardiac rehabilitation: long-term effects of a self-management intervention. *Soc Sci Med* 2006;62(12):3109-20

- Schrader G, Cheok F, Hordacre AL, et al. Predictors of depression three months after cardiac hospitalization. *Psychosom Med* 2004;66(4):514-20
- Schrader G, Cheok F, Hordacre AL, et al. Predictors of depression 12 months after cardiac hospitalization: the Identifying Depression as a Comorbid Condition study. *Aust N Z J Psychiatry* 2006;40(11-12):1025-30
- Schuit AJ, et al. Lichamelijke activiteit in samenhang met sociaal-demografische determinanten en gezondheidskenmerken bij volwassen mannen en vrouwen in Amsterdam, Doetinchem en Maastricht. *Ned Tijdsch Geneesk* 2001;143: 1559-64
- Schuler G, Hambrecht R, Schlierf G, et al. Regular physical exercise and low-fat diet. Effects on progression of coronary artery disease. *Circulation* 1992;86(1):1-11
- Schulte MB, Pluym B, Van Schendel G. Reintegration with duos: a self care program following myocardial infarction. *Patient Educ Couns* 1986;8:233-44
- Schwarz KA, Elman CS. Identification of factors predictive of hospital readmissions for patients with heart failure. *Heart Lung* 2003;32(2):88-99
- Scottish Intercollegiate Guidelines Network. Secondary prevention of coronary heart diseases following myocardial infarction. A National Clinical guideline. Scottish Intercollegiate Guidelines Network 2000
- Sears SF Jr, et al. Examining the psychosocial impact of implantable cardioverter defibrillators: a literature review. *Clin Cardiol* 1999;22(7):481-9
- Sebern M, Riegel B. Contributions of supportive relationships to heart failure self-care. *Eur J Cardiovasc Nurs* 2009;8(2):97-104
- Sebrechts CP, Klein JL, Ahnve S, et al. Myocardial perfusion changes following 1 year of exercise training assessed by thallium-201 circumferential count profiles. *Am Heart J* 1986;112(6):1217-26
- Sebregts EH, et al. Psychological effects of a short behavior modification program in patients with acute myocardial infarction or coronary artery bypass grafting. A randomized controlled trial. *J Psychosom Res* 2005;58(5):417-24
- Shaper AG, Wannamethee SG. Alcohol intake and mortality in middle aged men with diagnosed coronary heart disease. *Heart* 2000;83:394-9
- Shibeshi WA, Young-Xu Y, Blatt CM. Anxiety worsens prognosis in patients with coronary artery disease. *J Am Coll Cardiol* 2007;49(20):2021-7
- SHPIC (Scottish Health Purchasing Information Centre). Cardiac Rehabilitation SHPIC 1998 ([www.nhsconfed.net/scotland/shpic](http://www.nhsconfed.net/scotland/shpic))
- Shrey DE, Mital A. Accelerating the return to work (RTW) chances of coronary heart disease (CHD) patients: part 2--development and validation of a vocational rehabilitation programme. *Disabil Rehabil* 2000;10-20;22(13-14):621-6
- Simpson SH, et al. Using focus groups to identify barriers to drug use in patients with congestive heart failure. *Pharmacotherapy* 2000;20(7):823-9



- Singh RB, Niaz MA, Sharma JP, et al. Randomized, double-blind, placebo-controlled trial of fish oil and mustard oil in patients with suspected acute myocardial infarction: the Indian experiment of infarct survival-4. *Cardiovasc Drugs Ther* 1997;11:485-91
- Singh RB, Rastogi SS, Verma R, et al. An Indian experiment with nutritional modulation in acute myocardial infarction. *Am J Cardiol* 1992;69:879-85
- Sirois BC, Sears SF Jr, Bertolet B. Biomedical and psychosocial predictors of anginal frequency in patients following angioplasty with and without coronary stenting. *J Behav Med* 2003;26(6):535-51
- Sirois, B.C., S.F. Sears, Jr., and B. Bertolet, Biomedical and psychosocial predictors of anginal frequency in patients following angioplasty with and without coronary stenting. *J Behav Med* 2003;26(6):535-51
- Siscovick D, Weiss N, Fletcher R. The incidence of primary cardiac arrest during vigorous exercise. *N Engl J Med* 1984;311:874-77
- Slebus FG, Kuijer PPF, Willems JHB, et al. Prognostic factors for work ability in sicklisted employees with chronic diseases. *Occ Environ Med* 2007;64(12):814-9
- Smeulders ES, et al. The impact of a self-management group programme on health behaviour and healthcare utilization among congestive heart failure patients. *Eur J Heart Fail* 2009;11(6):609-16
- Smith OR, et al. Vital exhaustion in chronic heart failure: symptom profiles and clinical outcome. *J Psychosom Res* 2009;66(3):195-201
- Smith-Warner SA, Spiegelman D, Yaun SS, et al. Alcohol and breast cancer in women: a pooled analysis of cohort studies. *JAMA* 1998;279:535-40
- Snel, J. Alcohol, nuchter bekeken. Positieve effecten van matig gebruik. Assen: Van Gorcum 2002
- Soderman E, Lisspers J, Sundin O. Depression as a predictor of return to work in patients with coronary artery disease. *Soc Sci Med* 2003;56(1):193-202
- Soejima Y, Steptoe A, Nozoe S, et al. Psychosocial and clinical factors predicting resumption of work following acute myocardial infarction in Japanese men. *Int J Cardiol* 1999;72(1):39-47
- Sola CL, Bostwick JM. Implantable cardioverter-defibrillators, induced anxiety, and quality of life. *Mayo Clin Proc* 2005; 80(2):232-37
- Sørensen C, et al. Depression assessed over 1-year survival in patients with myocardial infarction. *Acta Psychiatr Scand* 2006;113(4):290-7
- Sørensen C, Friis-Hasché E, Haghfelt T, et al. Postmyocardial Infarction Mortality in Relation to Depression: A Systematic Critical Review. *Psychother Psychosom* 2005;74(2):69-80
- Southard BH, Southard DR, Nuckolls J. Clinical trial of an internet-based case management system for secondary prevention of heart disease. *J Cardiopulm Rehabil* 2003;23(5):341-8.

- Sparrow D, Dawbert TR. The influence of cigarette smoking on prognosis after a first myocardial infarction. A report from the Framingham study. *J Chronic Dis* 1978;31:425-32
- Spijkerman T, et al. Depression following myocardial infarction: first-ever versus ongoing and recurrent episodes. *Gen Hosp Psychiatry* 2005b;27(6): 411-17
- Spijkerman TA, et al. Who is at risk of post-MI depressive symptoms? *J Psychosom Res* 2005a; 58(5): 425-32; 433-34
- Spindler H, et al. Gender differences in anxiety and concerns about the cardioverter defibrillator. *Pacing Clin Electrophysiol* 2009;32(5): 614-21
- Stahle A, et al. Improved physical fitness and quality of life following training of elderly patients after acute coronary events. A 1 year follow-up randomized controlled study. *Eur Heart J*, 1999b. 20(20): 1475-84.
- Stahle A, Nordlander R, Ryden L, et al. Effects of organized aerobic group training in elderly patients discharged after an acute coronary syndrome. A randomized controlled study. *Scand J Rehabil Med* 1999a;31:101-7
- Steenland K, Hu S, Walker J. All-cause and cause-specific mortality by socioeconomic status among employed persons in 27 US states, 1984-1997. *Am J Public Health* 2004;94(6):1037-42
- Stern MJ, Gorman PA, Kaslow L. The group counseling exercise therapy study. A controlled intervention with subjects following myocardial infarction. *Arch Intern Med* 1983;143(9):1719-25
- Stevens VJ, Obarzanek E, Cook NR, et al. Long-term weight loss and changes in blood pressure: results of the Trials of Hypertension Prevention, phase II. *Ann Intern Med* 2001;134:1-11
- Stewart M, et al. Myocardial infarction: survivors' and spouses' stress, coping, and support. *J Adv Nurs* 2000;31(6):1351-60
- Straus SM, et al. The incidence of sudden cardiac death in the general population. *J Clin Epidemiol* 2004;57(1):98-102
- Strik JJ, Denollet J, Lousberg R, et al. Comparing symptoms of depression and anxiety as predictors of cardiac events and increased health care consumption after myocardial infarction. *J Am Coll Cardiol* 2003;42(10):1801-7
- Strik JJ, et al. Efficacy and safety of fluoxetine in the treatment of patients with major depression after first myocardial infarction: findings from a double-blind, placebo-controlled trial. *Psychosom Med* 2000;62(6):783-9
- Strik JJ, et al. One year cumulative incidence of depression following myocardial infarction and impact on cardiac outcome. *J Psychosom Res* 2004;56(1):59-66
- Strik JJ, van Praag HM, Honig A. Depression after first myocardial infarction. A prospective study on incidence, prognosis, risk factors and treatment. *Tijdschr Gerontol Geriatr* 2003;34(3):104-12

- Sullivan MJ, Green HJ, Cobb FR. Skeletal muscle biochemistry and histology in ambulatory patients with long-term heart failure. *Circulation* 1990;81(2):518-27
- Sullivan MJ, Higginbotham MB, Cobb FR. Exercise training in patients with chronic heart failure delays ventilatory anaerobic threshold and improves submaximal exercise performance. *Circulation* 1989;79(2):324-29
- Swan L, Hillis WS. Exercise prescription in adults with congenital heart disease: a long way to go. *Heart* 2000;83(6):685-7
- Swedeheart (Zweedse Hartstichting). Årsrapport 2008 (Jaarverslag 2008). Uppsala, Uppsala Clinical Research Center 2009.
- Swenson, J.R., S. Doucette, and D. Fergusson, Adverse cardiovascular events in antidepressant trials involving high-risk patients: a systematic review of randomized trials. *Can J Psychiatry*, 2006. 51(14): 923-9.
- Sykes DH, Hanley M, Boyle DM, Higginson JDS. Work strain and the post-discharge adjustment of patients following a heart attack. *Psychology & Health*. 2000;15(5):609-23.
- Tang JL, Armitage JM, Lancaster T, et al. Systematic review of dietary intervention trials to lower blood total cholesterol in free-living subjects. *BMJ* 1998;316:1213-20
- Taylor CB, et al. Effects of antidepressant medication on morbidity and mortality in depressed patients after myocardial infarction. *Arch Gen Psychiatry* 2005;62(7):792-8
- Taylor CB, Houston-Miller N, Ahn DK, et al. The effects of exercise training programs on psychosocial improvement in uncomplicated post-myocardial infarction patients. *J Psychosom Res* 1986;30(5):581-7
- Taylor CB, Houston-Miller N, Haskell WL, et al. Smoking cessation after acute myocardial infarction: the effects of exercise training. *Addict Behav* 1988;13(4):331-5
- Taylor RS, Watt A, Dalal HM, et al. Home-based cardiac rehabilitation versus hospital-based rehabilitation: a cost effectiveness analysis. *Int J Cardiol* 2007;119(2):196-201.
- Theobald K. The experience of spouses whose partners have suffered a myocardial infarction: a phenomenological study. *J Adv Nurs* 1997;26(3):595-601
- Theorell T, Oxenstierna G, Westerlund H et al. Downsizing of staff is associated with lowered medically certified sick leave in female employees. *Occup Environ Med* 2003;60(9):E9
- Thomas SA, et al. Quality of life and psychological status of patients with implantable cardioverter defibrillators. *Am J Crit Care* 2006; 5(4):389-98
- Thombs BD, Bass EB, Ford DE, et al. Prevalence of depression in survivors of acute myocardial infarction. *J Gen Intern Med* 2006;21(1):30-8
- Thombs BD, et al. Depression screening and patient outcomes in cardiovascular care: a systematic review. *JAMA* 2008b;300(18):2161-71
- Thombs BD, Ziegelstein RC, Whooley MA. Optimizing detection of major depression among patients with coronary artery disease using the patient health questionnaire: data from the heart and soul study. *J Gen Intern Med* 2008a;23(12):2014-17

- Thompson DR, Bowman GS. Evidence for the effectiveness of cardiac rehabilitation. *Intensive Crit Care Nurs* 1998;14(1):38-48
- Thompson DR, Meddis R. A prospective evaluation of in-hospital counseling for first time myocardial infarction men. *J Psychosom Res* 1990;34(3):237-48
- Thompson DR. A randomized controlled trial of in-hospital nursing support for first time myocardial infarction patients and their partners: effect on anxiety and depression. *J Adv Nurs* 1989;14(4):291-7
- Thompson PD, Buchner D, Pina IL, et al. Exercise and physical activity in the prevention and treatment of atherosclerotic cardiovascular disease : a statement from the Council on Nutrition, Physical Activity, and Metabolism (Subcommittee on Physical Activity). *Circulation* 2003;107:3109-16.
- Thun MJ, Peto R, Lopez AD, et al. Alcohol consumption and mortality among middle-aged and elderly U.S. adults. *N Engl J Med* 1997;337:1705-14
- Tiedtke C. Sociale participatie na een myocardinfarct: Exploratie van omvang en aard [Master Thesis]. Maastricht: Maastricht University 2007
- Todaro JF, Shen BJ, Raffa SD, et al. Prevalence of anxiety disorders in men and women with established coronary heart disease. *J Cardiopulm Rehabil Prev* 2007;27(2):86-91
- TOHP research group (Trails of Hypertension Prevention Collaborative research group). Effects of weight loss and sodium reduction intervention on blood pressure and hypertension incidence in overweight people with high-normal blood pressure. *Trails of Hypertension Prevention, Phase II. Arch Intern Med* 1997;157:657-67
- Truswell AS. Cereal grains and coronary heart disease. *Eur J Clin Nutr* 2002;56:1-14
- Truswell AS. Meta-analysis of the cholesterol-lowering effects of dietary fiber. *Am J Clin Nutr* 1999;70:942-3
- Truswell AS. Review of dietary intervention studies: effect on coronary events and on total mortality. *Aust N Z J Med* 1994;24:98-106
- Trzcieniecka-Green A, Steptoe A. The effects of stress management on the quality of life of patients following acute myocardial infarction or coronary bypass surgery. *Eur Heart J* 1996;17:1663-70
- Turner DA, Paul S, Stone MA, et al. Cost-effectiveness of a disease management programme for secondary prevention of coronary heart disease and heartfailure in primary care. *Heart* 2008;94(12):1601-6
- Tyni-Lenne R, Dencker K, Gordon A, et al. Comprehensive local muscle training increases aerobic working capacity and quality of life and decreases neurohormonal activation in patients with chronic heart failure. *Eur J Heart Fail* 2001;3:47-52
- Tyni-Lenne R, Gordon A, Jansson E, et al. Skeletal muscle endurance training improves peripheral oxidative capacity, exercise tolerance, and health-related quality of life in women with chronic congestive heart failure secondary to either ischemic cardiomyopathy or idiopathic dilated cardiomyopathy. *Am J Cardiol* 1997;80:1025-9

- Tyni-Lenne R, Gordon A, Jensen-Urstad M, et al. Aerobic training involving a minor muscle mass shows greater efficiency than training involving a major muscle mass in chronic heart failure patients. *J Card Fail* 1999;5:300-7
- Tyni-Lenne R, Gordon A, Sylven C. Improved quality of life in chronic heart failure patients following local endurance training with leg muscles. *J Card Fail* 1996;2:111-7
- Uniken Venema-van Uden MMAT. Hartrevalidatie. Over het bepalen en voorspellen van herstel. Proefschrift Erasmus Universiteit Rotterdam 1990
- USDHHS (U.S. Department of Health and Human Services). Cardiac Rehabilitation. Clinical practice guideline No. 17 (AHCOR publication No. 96-0672). U.S. Department for Health and Human Service. Agency for Health care Policy and Research 1995
- USDHHS (U.S. Department of Health and Human Services). The health benefits of smoking cessation. A Report of the Surgeon General. DHHS pub No (CDC)90-8416. Rockville, MD: US Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research USA and the National Heart, Lung, and Blood Institute 1990
- Utens EM, et al. Psychopathology in young adults with congenital heart disease. Follow-up results. *Eur Heart J* 1998;19(4):647-51
- Vaananen A, Koskinen A, Joensuu M, et al. Lack of predictability at work and risk of acute myocardial infarction: an 18-year prospective study of industrial employees. *Am J Public Health* 2008;98(12):2264-71
- Vahtera J, Kivimaki M, Pentti J, et al. Organisational downsizing, sickness absence, and mortality: 10-town prospective cohort study. *BMJ* 2004;328(7439):555
- Valliant PM, Leith B. Impact of relaxation training and cognitive therapy on coronary patients post surgery. *Psychol Rep* 1986;59:1271-78
- van Berkel TF, Boersma H, roos-Hesselink JW, et al. Impact of smoking cessation and smoking interventions in patients with coronary heart disease. *Eur Heart J* 1999;20:1773-82
- Van Camp SP, Peterson RA. Cardiovascular complication of out-patient cardiac rehabilitation programs. *JAMA* 1986;256:1160-3
- van den Broek KC, et al. Risk of ventricular arrhythmia after implantable defibrillator treatment in anxious type D patients. *J Am Coll Cardiol* 2009;54(6):531-37
- Van den Broek KC, et al. Shocks, personality, and anxiety in patients with an implantable defibrillator. *Pacing Clin Electrophysiol* 2008;31(7):850-57
- van der Meer, HCB. Taken en verantwoordelijkheden van de bedrijfsarts in het kader van de verzuimbegeleiding en re-integratie. Utrecht: Nederlandse Vereniging van Arbeids- en Bedrijfsartsen 2009
- Van Dijk JL, Bekedam, MA, Brouwer W, et al. Nederlandse Vereniging voor Arbeids- en Bedrijfsgeneeskunde Richtlijn Ischemische Hartziekten, in Handelen van de bedrijfsarts bij werknemers met ischemische hartziekten. Utrecht: NVAB Kwaliteitsbureau 2006.

- van Dixhoorn JJ, Duivenvoorden HJ, Pool J, et al. Psychic effects of physical training and relaxation therapy after myocardial infarction. *J Psychosom Res* 1990;34:327-37
- van Dixhoorn JJ, Duivenvoorden HJ, Staal JA, et al. Physical training and relaxation therapy in cardiac rehabilitation assessed through a composite criterion for training outcome. *Am Heart J* 1989;118:545-52
- van Dixhoorn JJ, Duivenvoorden HJ. Effect of relaxation therapy on cardiac events after myocardial infarction: a 5-year follow-up study. *J Cardiopulm Rehabil* 1999;19:178-85
- van Dixhoorn JJ, Significance of breathing awareness and exercise training for recovery after myocardial infarction. in: J.G. Carlson, A.R. Seifert, N. Birbaumer eds, *Clinical applied psychophysiology*, Plenum Press, New York, 1994, 113-132.
- van Dixhoorn JJ, White A. Relaxation therapy for rehabilitation and prevention in ischaemic heart disease: a systematic review and meta-analysis. *Eur J Cardiovasc Prev Rehabil* 2005; 12(3):193-202
- Van Dixhoorn, JJ. *Ontspanningsinstructie: principes en oefeningen*, Elsevier/Bunge: Maarssen 1998.
- van Elderen T, Dusseldorp E. Lifestyle effects of group health education for patients with coronary heart disease. *Psychology and Health* 2001;16:327-41
- van Elderen T, van der Kamp, LJT, Maes S, et al. Dutch translation of the 'Quality of Life after Myocardial Infarction' instrument. Leiden: vakgroep Gezondheidspsychologie, Universiteit Leiden 2000
- Van Engen-Verheul MM, Hellemans, I.M., Goud R et al. *Beslisboom Poliklinische Indiciestelling Hartrevalidatie 2010*. 2010, NVVC-CCPH – Commissie Cardiovasculaire Preventie en Hartrevalidatie, LMDO-H – Landelijk Multidisciplinair Overleg Hartrevalidatie 2010
- van Erp J, Hinnen C, Sanderman R. *Psychosociale zorg bij hart- en vaatziekten*. 's Gravenhage: Nederlandse Hartstichting 2006
- van Erp J, Schipper K. Effecten van psychosociale interventies bij hart- en vaatziekten: coronaire hartziekten, hartfalen en cerebrovasculaire aandoeningen. *Psychologie en Gezondheid* 2007;35:27-37
- van Gaal LF, Wauters MA, De Leeuw IH. The beneficial effects of modest weight loss on cardiovascular risk factors. *Int J Obes Relat Metab Disord* 1997;21:5-9
- van Heyningen C. Cholesterol lowering margarine may not be useful in healthy fat modified diet. *BMJ* 1999;319:186
- van Horn E, Fleury J, Moore S. Family interventions during the trajectory of recovery from cardiac event: an integrative literature review. *Heart Lung* 2002;31(3):186-98
- Van Ittersum M, de Greef M, van Gelder, et al. Fear of exercise and health-related quality of life in patients with an implantable cardioverter defibrillator. *Int J Rehabil Res* 2003;26(2):117-22

- van Melle JP, de Jonge P, Spijkerman TA, et al. Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis. *Psychosom Med* 2004;66(6):814-22
- van Melle JP, et al. Beta-blockers and depression after myocardial infarction: a multicenter prospective study. *J Am Coll Cardiol* 2006;48(11):2209-14
- van Melle JP, et al. Effects of antidepressant treatment following myocardial infarction. *Br J Psychiatry* 2007;190: 460-6
- van Melle JP, et al. Prediction of depressive disorder following myocardial infarction data from the Myocardial Infarction and Depression-Intervention Trial (MIND-IT). *Int J Cardiol* 2006;109(1):88-94
- van Melle JP, et al. Prognostic association of depression following myocardial infarction with mortality and cardiovascular events: a meta-analysis. *Psychosom Med* 2004;66(6): 814-22
- van Rijen EH, et al. Psychosocial functioning of the adult with congenital heart disease: a 20-33 years follow-up. *Eur Heart J* 2003;24(7):673-83
- van 't Veer P, Jansen MC, Klerk M, et al. Fruits and vegetables in the prevention of cancer and cardiovascular disease. *Public Health Nutr* 2000;3:103-7
- van Vegchel N, de Jonge J, Bosma H, et al. Reviewing the effort-reward imbalance model: drawing up the balance of 45 empirical studies. *Soc Sci Med* 2005;60(5):1117-31
- van Zyl LT, et al. Platelet and endothelial activity in comorbid major depression and coronary artery disease patients treated with citalopram: the Canadian Cardiac Randomized Evaluation of Antidepressant and Psychotherapy Efficacy Trial (CREATE) biomarker sub-study. *J Thromb Thrombolysis* 2009;27(1):48-56
- Varvaro FF. Family role and work adaptation in MI women. *Clin Nurs Res* 2000; 9(3):339-51
- Varvaro FF. Women with coronary heart disease: an application of Roy's Adaptation Model. *Cardiovasc Nurs* 1991;27(6):31-5
- Velicer WF, et al. Interactive versus noninteractive interventions and dose-response relationships for stage-matched smoking cessation programs in a managed care setting. *Health Psychol* 1999;8(1):21-8
- Velicer WF, Prochaska JO, Fava JL, et al. Interactive versus noninteractive interventions and dose-response relationships for stage-matched smoking cessation programs in a managed care setting. *Health Psychol* 1999;18(1):21-8
- Virtanen M, Kivimaki M, Elovainio M, et al. From insecure to secure employment: changes in work, health, health related behaviours, and sickness absence. *Occup Environ Med* 2003;60(12):948-53
- Voedingscentrum. Zo eet Nederland. Resultaten van de voedselconsumptiepeiling 1997-1998. 's Gravenhage: Voedingscentrum 1998

- Völler H, Hahmann H, Gohlke H, et al. Effect of inpatient rehabilitation on cardiovascular risk factors in patients with coronary heart disease (article in German). *Dtsch Med Wochenschr* 1999;124:817-23
- Völler H, Klein G, Gohlke H, et al. Secondary prevention in coronary patients after inpatient rehabilitation (article in German). *Dtsch Med Wochenschr* 2000;125:1457-61
- Vonder Muhll I, et al. Benefits of cardiac rehabilitation in the ninth decade of life in patients with coronary heart disease. *Am J Cardiol* 2002;90(6):645-48
- Voors AA, van Brussel BL, Plokker HM, et al. Smoking and cardiac events after venous coronary bypass surgery. A 15-year follow-up study. *Circulation* 1996;93:42-7
- Voors AA, van Brussel BL, Plokker HW, et al. Smoking and cardiac events after venous coronary bypass surgery. A 15-year follow-up study. *Circulation* 1996;93:42-47
- Voskuil DW, Feskens EJ, Katan MB, et al. Intake and sources of alpha-linolenic acid in Dutch elderly men. *Eur J Clin Nutr* 1996;50:784-87
- Vriend JWJ, Van der Velde ET, Mulder BJM. Adults with congenital heart disease: a growing population. *Neth Heart J* 2002;10:345-47
- Waalewijn RA, de Vos R, Koster RW. Out-of-hospital cardiac arrests in Amsterdam and its surrounding areas: results from the Amsterdam resuscitation study (ARREST) in 'Utstein' style. *Resuscitation* 1998;38(3):157-67
- Wachelder EM, et al. Life after survival: long-term daily functioning and quality of life after an out-of-hospital cardiac arrest. *Resuscitation* 2009;80(5):517-22
- Wallner S, Watzinger N, Lindschinger M, et al. Effects of intensified lifestyle modification on the need for further revascularization after coronary angioplasty. *Eur J Clin Invest* 1999;29(5):372-9
- Wallner S, Watzinger N, Lindschinger M, et al. Effects of intensified lifestyle modification on the need for further revascularization after coronary angioplasty. *Eur J Clin Invest* 1999;29:372-9
- Wang HX, Leineweber C, Kirkeeide R, et al. Psychosocial stress and atherosclerosis: family and work stress accelerate progression of coronary disease in women. The Stockholm Female Coronary Angiography Study. *J Intern Med* 2007;261(3):245-54
- Wang W, et al. Chinese couples' experiences during convalescence from a first heart attack: a focus group study. *J Adv Nurs* 2008;61(3):307-15
- Wannamethee SG, Shaper Ag, Walker M, et al. Physical activity and mortality in older men with diagnosed coronary heart disease. *Circulation* 2000;102(12):1358-63
- Watkins LL, Blumenthal JA, Davidson JR, et al. Phobic anxiety, depression, and risk of ventricular arrhythmias in patients with coronary heart disease. *Psychosom Med* 2006;68(5):651-6
- Webb GD, Williams RG. Care of the adult with congenital heart disease: introduction. *J Am Coll Cardiol* 2001;37(5):1166



- Webster RA, Thompson DR, Mayou RA. The experiences and needs of Gujarati Hindu patients and partners in the first month after a myocardial infarction. *Eur J Cardiovasc Nurs* 2002;1(1):69-76
- Welter J, van Litsenburg L, Kuijpers P, de Rijk A (Redactie) Adviesrapportage Psychosociale screening als onderdeel van de beslisboom Hartrevalidatie door de tijdelijke werkgroep Psychosociale screening. Interne rapportage vakgroep Sociale Geneeskunde, FHML, Maastricht University en MUMC+ / Cardiologie MUMC +, Maastricht 2010
- Welton NJ, et al. Mixed treatment comparison meta-analysis of complex interventions: psychological interventions in coronary heart disease. *Am J Epidemiol* 2009;169(9): 1158-65
- Wenger NK, Froelicher ES, Smith LK et al. Cardiac Rehabilitation. Clinical Practice Guideline No 17 (AHCPR Pub No. 96-0672). Rockville, MD: US Department of Health and Human Services, Public Health Service, Agency for Health Care Policy and Research USA and the National Heart, Lung, and Blood Institute 1995.
- White-Williams C, Jalowiec A, Grady K. Who returns to work after heart transplantation? *J Heart Lung Transplant* 2005;24(12):2255-61
- Whitlock EP, Orleans CT, Pender N, et al. Evaluating primary care behavioral counseling interventions: an evidence-based approach. *Am J Prev Med* 2002;22:267-84
- WHO (World Health Organisation). Obesity: preventing and managing the global epidemic. Report of a WHO Consultation (894-2000). WHO 2000
- Whooley M, et al. Editorial Interdisciplinary stepped care for depression after acute coronary syndrome. *Arch Int Med* 2010;170(7):585-6
- Wielenga RP, Huisveld IA, Bol E, et al. Exercise training in elderly patients with chronic heart failure. *Coron Artery Dis* 1998;9:765-70
- Wielenga RP, Huisveld IA, Bol E, et al. Safety and effects of physical training in chronic heart failure. Results of the Chronic Heart Failure and Graded Exercise study (CHANGE). *Eur Heart J* 1999;20:872-9
- Wilhelmsen L, Sanne H, Elmfeldt D, et al. A controlled trial of physical training after myocardial infarction: effects on risk factors, nonfatal reinfarction, and death. *Prev Med* 1975;4(4):491-508
- Wilhelmsson C, Vedin JA, Elmfeldt D, et al. Smoking and myocardial infarction. *Lancet* 1975;415-20.
- Wilk C, Turkoski B. Progressive muscle relaxation in cardiac rehabilitation: a pilot study. *Rehabil Nurs* 2001;26:238-42
- Willenheimer R, Erhardt L, Cline C, et al. Exercise training in heart failure improves quality of life and exercise capacity. *Eur Heart J* 1998;19:774-81
- Willenheimer R, et al. Effects on quality of life, symptoms and daily activity 6 months after termination of an exercise training programme in heart failure patients. *Int J Cardiol* 2001;77(1): 25-31

- Willich S, Lewis M, Lowel H. Physical exertion as a trigger of acute myocardial infarction. *N Engl J Med* 1993;329:1684-90
- Winterfeld H-J, Risch A, Siewert H, et al. Der Einfluss der konzentrativen Entspannung auf Blutdruck und Haemodynamik bei hypertonen Patienten mit koronaren Herzkrankheit und aortokoronarer Venenbypassoperation (ACVB). *Z Physioth* 1991;43:220-4
- Winterfeld H-J, Siewert H, Bohm J, et al. Autogenes Training bei hypertonen Regulationsstörungen nach aortokoronarer Venenbypass-Operation (ACVB) bei koronarer Herzkrankheit. *Innere Medizin* 1993;48:201-4
- Wojciechowski FL, et al. The relationship between depressive and vital exhaustion symptomatology post-myocardial infarction. *Acta Psychiatr Scand* 2000;102(5):359-65
- Wood D, De Backer G, Faergeman O, et al. Prevention of coronary heart disease in clinical practice: recommendations of the Second Joint Task Force of European and other Societies on Coronary Prevention. *Atherosclerosis* 1998;140:199-270
- Wood PD, et al. Increased exercise level and plasma lipoprotein concentrations: a one year randomized controlled study in sedentary middle-aged man. *Metabolism* 1983;32:31-9
- Wood PD, et al. The distribution of plasma lipoproteins in middle-aged male runners. *Metabolism* 1976;25:1249-57
- Wood PD, et al. The effect of exercise on plasma high density lipoproteins. *Lipids* 1979;14:417-27
- Worcester MC, Hare DL, Oliver RG, et al. Early programmes of high and low intensity exercise and quality of life after acute myocardial infarction. *BMJ* 1993;307(6914):1244-7
- Wosomu D, Bedford D, Ballantyne D. A comparison of the effects of strength and aerobic exercise training on exercise capacity and lipids after coronary artery bypass surgery. *Eur Heart J* 1996;17(6):854-63
- Xin X, He J, Frontini MG, et al. Effects of alcohol reduction on blood pressure: a meta-analysis of randomized controlled trials. *Hypertension* 2001;38:1112-7
- Yonezawa R, et al. Effects of phase II cardiac rehabilitation on job stress and health-related quality of life after return to work in middle-aged patients with acute myocardial infarction. *Int Heart J* 2009 ;50(3):279-90
- Yu CM, Lau CP, Chau J, et al. A short course of cardiac rehabilitation programme is highly cost effective in improving long-term quality of life in patients with recent myocardial infarction of percutaneous coronary intervention. *Arch Phys Med Rehabil* 2004;85(12):1915-22
- Yu DS, et al. Correlates of psychological distress in elderly patients with congestive heart failure. *J Psychosom Res* 2004;57(6):573-81
- Yu DS, et al. Living with chronic heart failure: a review of qualitative studies of older people. *J Adv Nurs* 2008;61(5):474-83

- Zamarra JW, Schneider RH, Besseghini I, et al. Usefulness of the Transcendental Meditation Program in the treatment of patients with coronary artery disease. *Am J Cardiol* 1996;77:867-70
- Zayac S, Finch N. Recipients' of implanted cardioverter-defibrillators actual and perceived adaptation: a review of the literature. *J Am Acad Nurse Pract* 2009;21(10): 549-56.
- Ziegelstein RC, et al. Can doctors and nurses recognize depression in patients hospitalized with an acute myocardial infarction in the absence of formal screening? *Psychosom Med* 2005;67(3):393-97
- Ziegelstein RC. Depression in patients recovering from a myocardial infarction. *JAMA* 2001;286(13):1621-7
- Zipes DP, Garson A. Jr. 26<sup>th</sup> Bethesda conference: recommendations for determining eligibility for competition in athletes with cardiovascular abnormalities. Task Force 6: arrhythmias. *J Am Coll Cardiol* 1994;24(4):892-99
- Zock PL, Kromhout D. Nutrition and health-fish fatty acids against fatal coronary heart disease. *Ned Tijdschr Geneesk* 2002;146:2229-33
- Zock PL, Urgert R, Hulshof PJ, et al. Transvetzuren in de voeding: risico op coronaire hartziekten. *Ned Tijdschr Geneesk* 1998;142:1701-4
- Zung Vu Tran, et al. Differential effects of exercise on serum lipid and lipoprotein levels seen with changes in body weight. *JAMA* 1985;254:919-24